



## Yield Stability of Newly Released Wheat Varieties in Multi-environments of Bangladesh

Md. Ashraful Alam<sup>1\*</sup>, Zahidul Islam Sarker<sup>1</sup>, Md. Farhad<sup>1</sup>, Md. Abdul Hakim<sup>1</sup>,  
Naresh Chandra Dev Barma<sup>2</sup>, Md. Israil Hossain<sup>3</sup>, Md. Mokhlesur Rahman<sup>4</sup>  
and Rabiul Islam<sup>5</sup>

<sup>1</sup>Wheat Research Centre (WRC), Bangladesh Agricultural Research Institute (BARI), Dinajpur, Bangladesh.

<sup>2</sup>Regional Wheat Research Centre (RWRC), Bangladesh Agricultural Research Institute (BARI), Joydebpur, Gazipur, Bangladesh.

<sup>3</sup>Regional Wheat Research Centre (RWRC), Bangladesh Agricultural Research Institute (BARI), Rajshahi, Bangladesh.

<sup>4</sup>Regional Agricultural Research Station (RARS), Bangladesh Agricultural Research Institute (BARI), Jamalpur, Bangladesh.

<sup>5</sup>Regional Agricultural Research Station (RARS), Bangladesh Agricultural Research Institute (BARI), Jessore, Bangladesh.

### Authors' contributions

This study was carried out in collaboration among the authors. All authors managed the literature searches, read and approved the final manuscript.

### Article Information

DOI: 10.9734/IJPSS/2015/14824

#### Editor(s):

- (1) Sławomir Borek, Department of Plant Physiology, Adam Mickiewicz University, Poland.  
(2) Marco Aurelio Cristancho, National Center for Coffee Research, Chinchiná, Caldas, Colombia.  
(3) A. Mujib, Department of Botany, Hamdard University, India.

#### Reviewers:

- (1) Anonymous, Bangladesh.  
(2) Azhar A. Kham, Bahauddin Zakariya University, College of Agriculture, Layyah, Pakistan.  
(3) Anonymous, China.

Complete Peer review History: <http://www.sciencedomain.org/review-history.php?iid=955&id=24&aid=8312>

Original Research Article

Received 23<sup>rd</sup> October 2014  
Accepted 17<sup>th</sup> February 2015  
Published 28<sup>th</sup> February 2015

### ABSTRACT

Varietal adaptability to environmental fluctuations is important for the stabilization of crop production on both region and season. In this context, one thousand six trials with four wheat varieties viz., BARI Gom 25, BARI Gom 26, BARI Gom 27 and BARI Gom 28 were conducted in forty districts of 8 agricultural regions in Bangladesh during 2013-14. Objectives of this study were to evaluate the mean yield performance and yield stability of the newly released varieties in

\*Corresponding author: E-mail: [ashrafulw@yahoo.com](mailto:ashrafulw@yahoo.com);

different regions of the country in different environmental status. Yield of 4 varieties significantly varied among the locations. The highest mean yield was obtained from Rangpur region (4275.15 kg ha<sup>-1</sup>) followed by Jessore region (4266 kg ha<sup>-1</sup>). Yield of these two regions were statistically similar but significantly higher than other regions. In AMMI analysis, AMMI-1 biplot showed that Rangpur and Rajshahi were favorable environments; Sylhet and Comilla were unfavorable environments. Considering varieties, BARI Gom 26 and BARI Gom 25 were more stable and can be recommended for all environmental condition of Bangladesh. On the other hand, BARI Gom 27 was the most sensitive to changing environments. In AMMI-2 biplot, Comilla, Sylhet and Dhaka were the most discriminating environments for all tested varieties, while BARI Gom 26 and BARI Gom 25 were more stable varieties. The most suitable varieties for Rangpur, Rajshahi and Mymensingh was BARI Gom 26; BARI Gom 25 for Sylhet; BARI Gom 27 for Comilla, Jessore and Barishal while BARI Gom 28 was suitable for Dhaka. Among the varieties, BARI Gom 26 and BARI Gom 25 had higher mean grain yield values coupled with small positive IPCA 1 scores and hence these are recommended for all the locations.

*Keywords: Yield stability; Triticum aestivum; adaptability; AMMI analysis.*

## ABBREVIATION

*PTOS- Power tiller operated seeder; BpLB-Bipolarities Leaf Blight, BARI- Bangladesh agricultural research institute; WRC- Wheat research centre; MET- Multi-environment trials; AMMI- Additive main effect and multiplication interaction; GGE- Genotype main effects and genotype × environment interaction effects; PCA- Principal component analysis; DAE- Department of agricultural extension; NGOs- Non-government organizations; OFRD- On-farm research division; CSISA- Cereal systems initiative for south Asia; ARS- Agricultural research station; SAAOs- Sub-assistant agricultural officers; SA- Scientific assistance; SSAs- Senior scientific assistances; UAOs- Upazila agriculture officers; CPS- Crop protection specialist; IPCA- Interaction principal component axes; GEI- Genotyp-by-environment interaction; ASV- AMMI stability value.*

## 1. INTRODUCTION

The agricultural sector plays a very significant role in the Bangladesh economy as well socially and culturally. Agriculture accounts for about 20% of the country's Gross Domestic Product (GDP) [1]. In addition, agriculture provides more than 60% of the national employment. The agriculture sector is an important tool to reach different goals stated in the Poverty Reduction Strategy as it directly reaches the poorest in the rural areas, providing both food and cash at the same time. Wheat is the second most important cereal crop after rice in Bangladesh [2] and its consumption is increasing 3% per year [3]. The tragic irony is that the wheat production in the country 1.4 million tons [4], which is much below than the annual requirement. This deficit accounts about 75% of country's annual consumption and it is met, through imports [5]. Wheat production has increased steadily from around 0.115 million tons in 1971-72 to 1.9 million tons in 1999 and then gradually decreased to 0.763 million tons in 2006-07 [6,7]. This low production is mainly due to the reduction of area under wheat cultivation. The

highest wheat growing area was 0.85 million hectares in 1999 which now came down to 0.45 (2013-14) million ha [8]. Wheat is to compete with other profitable crops like Boro rice, corn, potato and winter vegetables which insisting farmers to push wheat crop in marginal lands from fertile ones and this is considered as the main reason for decreasing area under wheat [9]. Yield was also decreasing due to cultivation of old varieties that are susceptible to leaf rust and BpLB, knowledge gap about recommended technologies [10].

For increasing wheat production in Bangladesh, It is very important to adopt new technologies recommended by Wheat Research Centre for increasing wheat production in Bangladesh [11]. Use of power tiller operated seeder (PTOS) to confirm timely seeding and use of wheat thresher is also important to save the quality of seeds from early monsoon. WRC of Bangladesh Agricultural Research Institute (BARI) has released 30 varieties in conventional breeding approach and most of the later released varieties are better than previous one [12] in respect of yield, disease and terminal heat tolerance. But those varieties and other technologies are not

being adopted by the farmers in a satisfactory rate due to their inadequate knowledge about the varieties and technologies; and insufficient extension efforts [13].

Crop adaptability is commonly understood as wide or local adaptation as a result of high mean grain yield and yield stability across environments and seasons. On the other hand, yield stability of a cultivar is defined as consistency of yield response of variety(ies) across environments or seasons [14]. Varietal adaptability to environmental fluctuations is important for the stabilization of crop production on both region and season. Information on the genotype x environment interaction is essential to the successful evaluation of stable genotypes [15]. Multi-environment trials (MET) are conducted to evaluate the yield stability performance of different genetic materials under different environmental conditions [15,16]. Oftentimes, genotypes subject to different environments show significant fluctuations on the yield responses. These changes are influenced by different environmental conditions commonly referred to as genotype-by-environment (GE) interaction [17]. The action of GE interaction makes genetic progress difficult since it minimizes the association of phenotypic and genotypic values [18]. Hence, GE interaction should be exploited through selection of superior genotypes on specific location or avoided by selecting genotypes with wide adaptability and stable yield response across diverse locations [19]. Recently, several methods in analyzing GE interaction and MET data to reveal patterns of GE have been reported. These methods include regression coefficient [20], sum of squared deviations from regressions [21], stability of variance [22], coefficient of determination [23] coefficient of variability [24], and multiplicative main effects and multiplicative interaction (AMMI) [25,26,27] which have been commonly used. Another method proposed by Yan et al. [15] called GGE biplot analysis for graphical display of GE interaction pattern in MET data. GGE biplot is an effective method based on principal component analysis (PCA) which graphically displays GE in a two-way table and allows visual examination of the relationship interactions among test environments, genotypes, and GE interaction. The main objective of this study was to evaluate the yield performance and yield stability of the newly released varieties in different regions of the country in different environmental status.

## 2. MATERIALS AND METHODS

One thousand six trials were conducted with 4 newly released wheat varieties viz., BARI Gom 25 & BARI Gom 26 (released 2010) and BARI Gom 27 & BARI Gom 28 (released 2012) in the farmers' fields of forty districts out of sixty four under 8 agricultural regions in 2013-14 (Table 1 & Fig. 1) in collaboration with Department of Agricultural Extension (DAE), Non-government Organizations (NGOs), Cereal Systems Initiative for South Asia (CSISA), On Farm Research Division (OFRD), & Agricultural Research Station (ARS) of BARI. Seven hundred twenty three trials were conducted in collaboration with DAE, eighty through NGOs, forty through CSISA and one hundred fifty one trials by BARI scientists (thirty six through WRC & ARS and one hundred fifteen through OFRD). The pedigree of these varieties are BARI Gom 25 (ZSH 12/HLB 19//2\*NL 297), BARI Gom 26 (ICTAL123/3/RAWAL87//VEE/HD2285), BARI Gom 27 (FRANCOLIN # 1) and BARI Gom 28 (CHIL/2\*STAR/4/BOW/CROW//BUC/PVN/3/2\*VE E#10). Each trial had a single variety with a plot size of 800 m<sup>2</sup>. with no replication. Each of the variety was allotted to a single farmer. Only seed, fertilizers, signboard and printed documents on wheat production technologies were supplied to farmers. All other management practices were done by farmers. A day long training was given to the farmers, related Sub Assistant Agricultural Officer (SAAOs), Scientific Assistance (SA)/Senior Scientific Assistances (SSAs), Upazila Agriculture Officers (UAOs) and Crop Protection Specialist (CPS) of the respective districts about trials, wheat seed production and preservation technologies. Seeds were sown from 10 November to 16 December 2013. Weekly temperature data recorded during the growing period of wheat is graphically presented in Figs. 2a & 2b.

Yields were collected from 20 m<sup>2</sup> (4 samples×5 m<sup>2</sup>) areas of each trial. A data collection template was supplied to farmers for sending data on yield, sowing & harvesting date, number of irrigations, seed preservation, overall comments, etc. through supervising officials of DAE, NGOs and BARI.

In AMMI model, the contribution of each genotype and each environment to the GEI is assessed by the use of the biplot graph display where yield means are plotted against the scores of the IPCA1 [27]. The stability parameters and regression coefficient (bi) were estimated

according to Eberhart and Russell [21]. All the data were subjected to analysis using two software programmes: Cropstat version 7.2 after Zobel et al. [27] and SAS. 9.

The AMMI model does not make provision for a quantitative stability measure, such a measure is essential in order to quantify and rank genotypes

according to their yield stability, the following measure was proposed by Purchase [28]:

$$ASV = \sqrt{\frac{[IPCA1 \text{ sum of square } (IPCA1 \text{ score})]^2}{[IPCA2 \text{ sum of square } + [IPCA2 \text{ score}]^2}}$$







Fig. 1. Showing trials located in different regions of Bangladesh




Weekly maximum, minimum and average temperature in Rangpur region



Weekly maximum, minimum and average temperature in Rajshahi region




Weekly maximum, minimum and average temperature in Mymensingh region




Weekly maximum, minimum and average temperature in Jessore region


Fig. 2a. Weather data containing minimum, maximum and average temperature of Rangpur, Rajshahi, Jessore and Mymensingh region during the experimental season




Weekly maximum, minimum and average temperature in Dhaka region



Weekly maximum, minimum and average temperature in Comilla region



Weekly maximum, minimum and average temperature in Barisal region



Weekly maximum, minimum and average temperature in Sylhet region

Fig. 2b. Weather data containing minimum, maximum and average temperature of Dhaka, Comilla, Sylhet and Barisal region during the experimental season

### 3. RESULTS AND DISCUSSION

#### 3.1 Agronomic Performance

Yield of 4 varieties significantly ( $F_{(8,20)} = 8.65$ ;  $P = < 0.001$ ) varied among the locations. The highest mean yield was obtained from Rangpur region ( $4275.15 \text{ kg ha}^{-1}$ ) followed by Jessore region ( $4266 \text{ kg ha}^{-1}$ ). Yield of these two regions were statistically similar but significantly higher than other regions (Table 2). The lowest mean yield was recorded in Sylhet region due to the water stress. Hasan et al. [29] found that area and production of wheat were fluctuating due to weather condition and technological changes based on secondary data during 1980/81-2003/04. This reference support current findings of variation in wheat yield among locations. The yield of BARI Gom 26 was highest ( $4725 \text{ kg ha}^{-1}$ ) in Rangpur region followed by BARI Gom 28 ( $4530 \text{ kg ha}^{-1}$ ) in Dhaka region and BARI Gom 25 ( $4370.70 \text{ kg ha}^{-1}$ ) in Jessore region. At Rajshahi region, the yield of BARI Gom 26 was highest ( $4302.29 \text{ kg ha}^{-1}$ ) followed by BARI Gom 25 ( $4006.71 \text{ kg ha}^{-1}$ ). The BARI Gom 27 produced lowest yield ( $27.20 \text{ kg ha}^{-1}$ ) in Sylhet region. The highest mean yield was recorded in BARI Gom 26 ( $3977.83 \text{ kg ha}^{-1}$ ) followed by BARI Gom 28 ( $3828.01 \text{ kg ha}^{-1}$ ), BARI Gom 25 ( $3819.31 \text{ kg ha}^{-1}$ ) and BARI Gom 27 ( $3709.78 \text{ kg ha}^{-1}$ ). BARI Gom 26 produced the highest yield but statistically ( $F_{(3,20)} = 1.42$ ;  $P = 0.265$ ) similar with all other varieties. Mohammed [30] observed in Northern Sudan using AMMI analysis that considering the average performance combined over environments, the genotypes differed significantly in grain yield of bread wheat. Naroui Rad et al. [31] also found variation in grain yield of bread wheat lines due to environmental variation.

The farmer have preserved total 95426 kg seeds of the new varieties. The highest seed stored ( $27925 \text{ kg}$ ) in Jessore region followed by

Rajshahi region ( $23423 \text{ kg}$ ). In total 50050 farmers of the same and neighboring villages visited the trials and expressed their interest to collect seeds of new varieties for the next year. These results demonstrated that all of 4 new varieties (BARI Gom 25, BARI Gom 26, BARI Gom 27 and BARI Gom 28) were preferred by the farmers.

#### 3.2 Stability Performance

Adaptability of the new wheat varieties over wide ranges of environments (40 districts) are presented in Fig. 3 which indicates that BARI Gom 25 and BARI Gom 26 were more stable than other varieties. These varieties had comparatively higher yields in poor environment and can be recommended for growing in poor environments. BARI Gom 27 was more sensitive to changing production environments. It had low yield in poor and very high yields in better environments. This variety could be recommended for growing in favourable environments. BARI Gom 28 had average sensitivity to changing production environments.

#### 3.3 AMMI Biplot Analysis

Table 3 reflects IPCA score and AMMI stability value (ASV). BARI Gom 28 showed the lowest scores in the IPCA 1, followed by BARI Gom 26 and BARI Gom 25. BARI Gom 27 scored the highest IPCA 1 value. Considering The AMMI stability value (ASV) that takes into account the scores of the IPCA 2, the variety BARI Gom 28 appeared to be among those showing low ASV. The variety BARI Gom 26 became the second lowest in ASV. With regards to locations, Rangpur and Jessore gave the lowest IPCA 1 scores whereas Mymensingh, Rajshahi and Dhaka gave medium score and Comilla scored the highest IPCA 1 value.

**Table 1. Regional and Institutional distribution of wheat trials in the year 2013-14**

SI #	Region	Number of trials					Total
		DAE	NGO	WRC & ARS	CSISA	OFRD	
1	Rangpur	204	36	12	0	0	252
2	Rajshahi	190	36	12	0	0	238
3	Jessore	116	0	0	0	24	140
4	Mymensingh	72	0	12	36	12	132
5	Dhaka	24	0	0	0	15	39
6	Comilla	12	0	0	0	12	24
7	Sylhet	12	0	0	0	12	24
8	Barisal	93	12	0	12	40	157
Total		723	84	36	48	115	1006

**Table 2. Region-wise results of variety trials conducted through all organization (DAE, OFRD, ARS, WRC, CSISA & NGOs) in 2013-14**

Districts	No. of trials	Result of trials	Grain yield (kg/ha)				Mean yield ± SE	Seeds preserved by farmers (kg)
			BARI Gom25	BARI Gom26	BARI Gom27	BARI Gom28		
Rangpur	252	98	4002.50	4725.00	4116.50	4256.60	4275.15±4.45 a	17281
Rajshahi	238	118	4006.71	4302.29	3739.40	3895.71	3986.03±3.85 abc	23423
Jessore	140	140	4370.70	4212.80	4157.00	4324.40	4266.23±2.48 a	27925
Mymensingh	132	132	3692.88	3755.25	3363.63	3703.63	3628.84±3.34 bdc	11780
Dhaka	39	39	3983.33	4180.00	4050.00	4530.00	4185.83±3.90 ab	4770
Comilla	24	19	3290.00	3445.00	3840.00	3325.00	3475.00±3.97 cd	1807
Sylhet	24	10	3715.00	3640.00	2720.00	2950.00	3256.25±5.57 d	1170
Barishal	157	93	3493.33	3562.33	3691.67	3638.75	3596.52±2.33 bdc	7270
Total	1006	649						95426
Mean			3819.31	3977.83	3709.78	3828.01		

**Table 3. IPCA axes scores for genotypes and environments, AMMI stability value (AVS) and mean performance for grain yield (kg/ha) of 4 wheat varieties grown at 8 agricultural regions in Bangladesh (2013-14)**

Variety	Yield	Rank	IPCA1 Score	IPCA2 Score	bi	ASV	Rank	Location	IPCA1 Score
BARI Gom 25	3819.31	3	16.71	8.955	0.697	57.7	3	Rangpur	0.52
BARI Gom 26	3977.83	1	13.01	-4.08	0.994	44.6	2	Rajshahi	6.39
BARI Gom 27	3709.78	4	-20.75	11.89	1.007	71.8	4	Jessore	-1.52
BARI Gom 28	3828.01	2	-8.959	-16.76	1.303	34.9	1	Mymensingh	3.43
								Dhaka	-8.32
								Comilla	-14.56
								Sylhet	23.22
								Barishal	-9.15

The AMMI1 biplot provide a graphical representation of the relationship between the first interaction principal component axis (AMMI component 1) and mean of varieties and location [32] with the biplot according for up to 91.5% of the treatment sum of squares (Fig. 4). Variety and locations on the same parallel line, relative to ordinate, have similar yields and a variety or location on the right side of the midpoint of this axis has higher yields than those on the left hand side [27]. According to AMMI-1 biplot analysis (Fig. 4), environments showed high variation in both main effects and interactions (IPCA1). While the Rangpur and Jessore had low interaction; Sylhet and Comilla were highly interactive. Rajshahi was the most favorable environment and Comilla and Barishal was the least favorable environments, while Mymensingh and Dhaka were the average environments. Environment Rangpur and Jessore with IPCA1 scores near zero had no interaction effects, Variety BARI Gom 26 and location Rangpur combinations with IPCA1 scores of the same sign produced positive specific interaction effects and BARI Gom 28 and location Jessore combinations with IPCA1 score produced positive specific interaction effects. Hagos and Abay [33] also found the significant variation among genotype, location and genotype by location interaction in bread wheat by AMMI biplot analysis.

Consequently, the variety BARI Gom 26 was the high yielder followed by BARI Gom 28, BARI Gom 25 and BARI Gom 27. The AMMI1 estimation had a profound effect in producing clear and stratified ranking patterns and on this basis BARI Gom 26 and BARI Gom 25 would be

considered more adapted to a wide range of environments than the rest of the varieties. BARI Gom 27 had low yield and unstable and BARI Gom 28 had high yield but moderately stable. Dixon and Nukenine [34] and Crossa et al. [35] obtained a similar stratification in cassava and wheat genotypes, respectively.

According to AMMI-2 biplot analysis (Fig. 5), the varieties (and environments) that are located far away from the origin are more responsive. BARI Gom 27 BARI Gom 28 appeared unstable due to their dispersed position. Comilla, Sylhet and Dhaka were the most differentiating environments, while BARI Gom 27 and BARI Gom 28 were the most responsive genotypes. BARI Gom 26 and BARI Gom 25 appeared to be more stable when plotting the IPCA1 and IPCA2 scores. Genotypes and environments that fall in the same sectors interact positively; in contrast if they fall in opposite sectors interact negatively. If they fall into adjacent sectors, interaction is somewhat more complex. In this case, the best variety with respect to sites Dhaka was BARI Gom 28 but unsuitable for Sylhet. Variety BARI Gom 25 was suitable for Sylhet but unsuitable for Dhaka. With respect to Comilla, Barishal and Dhaka, variety BARI Gom 27 was particularly suitable but unsuitable for Rajshahi and Mymensingh, while variety BARI Gom 26 was well fit for Rajshahi and Mymensingh. For multivariate approach, the AMMI model is better for partitioning the G x E into the causes of variation, which ease identification of environments' potential and is used to identify superior genotypes either with specific adaptation or wide adaptation [32,36,37].






Fig. 3. Adaptability of the new wheat varieties conducted by DAE



**Fig. 4. AMMI 1 model for grain yield (kg ha<sup>-1</sup>) showing the means of varieties and locations against their respective IPCA 1 scores. RN = Rangpur, RJ= Rajshahi, JE= Jessore, MY= Mymensingh, DH= Dhaka, CO= Comilla, SY= Sylhet and BA= Barishal, 1 = BARI Gom 25, 2 = BARI Gom 26, 3 = BARI Gom 27, 4 = BARI Gom 28, Δ = Variety combination (genotypes), ● = Locations (8)**



**Fig. 5. AMMI 2 model for grain yield (kg ha<sup>-1</sup>) showing the IPCA scores of wheat varieties planted across locations**

#### 4. CONCLUSION

The AMMI statistical model has been used to diagnose the GE interaction pattern of grain yield of newly released wheat varieties. The most suitable varieties for Rangpur, Rajshahi and Mymensingh were BARI Gom 26; BARI Gom 25 for Sylhet; BARI Gom 27 for Comilla, Jessore and Barishal while variety BARI Gom 28 was suitable for Dhaka. BARI Gom 26 and BARI Gom 25 were hardly affected by the GE interaction and thus will perform well across a wide range of environments and these can be recommended for all locations. Locations, such as Rangpur and Rajshahi, could be regarded as a good selection sites for wheat cultivation for its stable yielding environment.

#### ACKNOWLEDGEMENTS

The authors are highly thankful to the farmers who were actively involved in trial management in their own lands. The author also acknowledge to the DAE and NGO representatives who have legitimate contribution to the trial management in the remote locaton. This experiment was carried out by Government fund of BARI.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. NAPA (National Adaptation Programme of Action). Climatic Risks Assessment at Community Level in Agriculture; 2012.
2. Sarker MAZ, Alam MA, Hossain A, Mannaf MA. Agro-economic performance of crop diversification in rice based cropping systems of northwest Bangladesh. Agriculture, Forestry and Fisheries. 2014;3(4):264-270.
3. Karim MA, Awal MA, Akter M. Forecasting of wheat production in Bangladesh. Bangladesh J Agril Res. 2010;35(1):17-28.
4. USDA. Bangladesh wheat production by year; 2014. Available:<http://www.indexmundi.com/agriculture/?country=bd&commodity=wheat&graph=production>
5. Hussain SS. Bangladesh grain and feed annual. USDA foreign agricultural service BARI. Year of release and average yield of Bangladesh wheat varieties develop since 1974; 2012.
6. BBS (Bangladesh Bureau of Statistics). Statistical year of Bangladesh; 2007.
7. Hossain A, Teixeira da Silva JA. Phenology, growth and yield of three wheat (*Triticum aestivum*) varieties as affected by high temperature stress. Not. Sci. Biol. 2013;4(3):97-109.
8. BBS (Bangladesh Bureau of Statistics). Statistical year of Bangladesh; 2014.
9. WRC (Wheat Research Centre). Annual Report, 2008-09. 2009;181.
10. Sufian MA. Sustainable wheat production in Bangladesh in relation to climate change. A Key note paper presented at the workshop on "Sustainable Wheat Production, held at BRAC Centre Inn, Dhaka; 2005.
11. WRC (Wheat Research Centre). Annual Report, 2013-14; 2014.
12. BARI (Bangladesh Agricultural Research Institute). BARI annual report 2012-13; 2013.
13. Pandit DB, Baksh ME, Sufian MA, Harunur-Rashid M, Islam MM. Impacts of participatory variety selection in wheat on agro-economic changes of wheat farmers in Bangladesh. Bangladesh J Agril Res. 2007;32(3):335-347.
14. Farshadfar E, Mohammadi R, Aghaee, M Vaisi, Z. GGE biplot analysis of genotype x environment interaction in wheat-barley disomic addition lines. Australian Journal of Crop Science. 2012;6(6):1074-1079.
15. Yan W, Hunt LA, Sheng Q, Szlavnic Z. Cultivar evaluation and mega-environment investigation based on the GGE biplot. Crop Science. 2000;40:597-605.
16. Delacy IH, Basford KE, Cooper M, Bull JK. Analysis of environment trials—an historical perspective. In: Cooper, M. and G.L. Hammer (Eds.). Plant Adaptation and Crop Improvement, CAB International, U.K.; 1996.
17. Allard RW. Bradshaw AD. Implication of genotype-environmental interaction in applied plant breeding. Crop Science. 1964;5:503-506.
18. Comstock RE, Moll RH. Genotype x Environment Interactions. Symposium on Statistical Genetics and Plant Breeding. National Academy Science Research Council, Washington, D.C. 1963;164-196.
19. Cecaelli S. Wide adaptation. How wide. Euphytica. 1989;40:197-205.
20. Finlay KW, Wilkinson GN. The analysis of adaptation in a plant breeding programme.

- Australian Journal of Agricultural Research. 1963;14:742-754.
21. Eberhart SA, Russel WA. Stability parameters for comparing varieties. *Crop Science*. 1966;6:36-40.
  22. Shukla GK. Some statistical aspects of partitioning of genotype-environmental components of variability. *Heredity*. 1972;29:237-245.
  23. Pinthus JM. Estimate of genotype value: A proposed method. *Euphytica*. 1973;22:121-123.
  24. Francis TR, Kannenberg LW. Yield stability studies in short season maize. I. A descriptive method for grouping genotypes. *Canadian Journal of Plant Science*. 1978;58:1029-1034.
  25. Gauch HG, RW Zobel. Predictive and postpredictive success of statistical analyses of yield trials. *Theoretical and Applied Genetics*. 1988;76:1-10.
  26. Gauch HG. *Statistical Analysis of Regional Yield Trials: AMMI Analysis of Factorial Designs*. Elsevier, Amsterdam, The Netherlands; 1992.
  27. Zobel RW, Wright MJ, Gauch HG. Statistical analysis of a yield trial. *Agronomy Journal* 1988;80:388-393.
  28. Purchase RL. Parametric analysis to describe genotype by environment interaction and yield stability in winter wheat. Ph.D. Thesis, Department of Agronomy, Faculty of Agriculture of the University of the Free State, Bloemfontein, South Africa; 1997.
  29. Hasan, MN, MA Monayem Miah, MS Islam QM Alam and MI Hossain. Change and instability in area and production of wheat and maize in Bangladesh. *Bangladesh J. Agril. Res*. 2008;33(3): 409-417.
  30. Maarouf I. Mohammed. Genotype X Environment Interaction in Bread Wheat in Northern Sudan Using AMMI Analysis *American-Eurasian J. Agric. & Environ. Sci*. 2009;6(4):427-433.
  31. Naroui Rad MR, Abdul Kadir M, Rafii MY, Hawa Z. E. Jaafar, Naghavi MR, Farzaneh Ahmadi. Genotype × environment interaction by AMMI and GGE biplot analysis in three consecutive generations of wheat (*Triticum aestivum*) under normal and drought stress conditions. *Australian Journal of Crop Science*. 2013;7(7):956-961.
  32. Kempton RA. The Use of Biplots in interpreting Variety by Environment Interactions. *Journal of Agricultural Science*. 1984;103:123-135.
  33. Hagos HG, Abay F. AMMI and biplot analysis of bread wheat genotypes in the northern part of Ethiopia. *J. Pl. Bred & Genet*. 2013;1(1):12-18
  34. Dixon AGO, Nukene EN. Statistical analysis of cassava yield trials with the additive main effects and multiplicative interaction (AMMI) model. *Afr J Root Tuber Crops*. 1997;3:46-50.
  35. Crossa J, Fox PN, Pfeiffer WH, Rajaram S, Gauch HG. AMMI adjustment for statistical analysis of an international wheat yield trial. *Theor Appl Genet*. 1991;81:27-37.
  36. Anandan A, Eswaran R, Sabesan T, Prakash M. Additive Main Effects and Multiplicative Interactions Analysis of Yield Performances in Rice Genotypes under Coastal Saline Environments. *Advances in Biological Research*. 2009;3(1-2):43-4.
  37. Crossa J. Statistical analyses of multi location trials. *Advances in Agronomy*. 1990;44:55.

© 2015 Alam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sciencedomain.org/review-history.php?iid=955&id=24&aid=8312>