



Insecticidal and repellent activities of *Azadirachta indica* A. (Juss) oil against fifth instar larvae and adults of *Sahlbergella singularis* (Hemiptera:Miridae)

**Koffi Gagnon Ankou^{a*}, Lankondjoa Kolani^a,
Essolakina Magnim Bokobana^a, Abouwaliou Nadio^a,
Koffi Koba^a and Komla Sanda^a**

^a *Laboratoire de Recherche sur les Agroressources et la Santé Environnementale (LARASE) Ecole Supérieure d'Agronomie, Université de Lomé, BP. 20131, Lomé, Togo.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAHR/2023/v10i4256

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101065>

Original Research Article

Received: 12/04/2023
Accepted: 14/06/2023
Published: 22/06/2023

ABSTRACT

Sahlbergella singularis is one of the insects that causes the most damage to cocoa trees in Togo. The management of this pest is mainly based on chemical control, with its negative consequences for human health and the environment. In order to find an alternative to the use of synthetic pesticides, the insecticidal and repellent activities of neem oil were tested in vitro on fifth instar larvae and adults of *S. singularis*. Larvae were collected from a cocoa plot. The fifth instar larvae were separated from the other stages. Some of the larvae were reared in the laboratory until adult

*Corresponding author: Email: ankoukoffi@gmail.com;

emergence. The concentrations of neem oil used were 0.5, 1, 1.5, 2 and 2.5 $\mu\text{l.ml}^{-1}$. Insecticidal tests were carried out by spraying the solutions directly onto the insects. The repellency test was carried out using the preferential zone method on filter paper. The results showed an increasing mortality rate with increasing concentrations of neem oil for both fifth instar larvae and adults. The calculated LC50's for larvae and adults were 0.89 and 0.98 $\mu\text{l.ml}^{-1}$ respectively. Neem oil had a weak repellent effect on both fifth instar larvae and adults, with average repellency rates of 23.60% and 28.8% respectively. These results show that neem oil can be an alternative to the use of synthetic pesticides in the control of *S. singularis*.

Keywords: *Azadirachta indica*; *Sahlbergella singularis*; larvae; adults; insecticidal; repellent.

1. INTRODUCTION

Originally from the Amazon, the cocoa tree is a plant widespread in the world and known as the food of the gods [1]. In several countries in general and Togo in particular, it is considered a cash crop [2]. Most of the world's cocoa production, 70%, comes from Africa [3] and occupies an important place in the economy of the producing countries in West Africa. In Togo, cocoa production occupies 15,414 households [4], and its contribution to Togo's GDP was estimated at 1.3% in 2010 [5]. National cocoa production is estimated at more than 10,000 t with a total export of 9,127 t in 2020 [6]. Cocoa production faces many constraints, including lack of maintenance of orchards, poor management of shade, poor management of soil fertility and old orchards, and especially poor management of insects, diseases, and pests [3].

Among these factors, diseases and insects cause the most loss. Indeed, the cocoa tree is a plant strongly attacked by insects and diseases. Mirids (*Distantiella theobroma* and *Sahlbergella singularis*) are the main insect pests of cocoa in West Africa in general and in Togo in particular. They are responsible for losses ranging from 20 to 75% [7,8]. There are several methods of controlling these insects, but chemical control is the most widely used [9].

It is known that the use of synthetic chemical pesticides pollutes groundwater, water sources, plants, and the ecosystem [10]. In cocoa production, the production and use of synthetic chemical fertilizers and pesticides are major causes of environmental pollution [11]. In addition, analysis of bean samples revealed the presence of organophosphates and synthetic pyrethroids [12,13]. These pesticides are thus ingested by consumers and are found in blood, serum, and sweat [14,15]. Exposure to synthetic chemical pesticides is associated with cancer of the blood, liver, prostate, pancreas, uterus, and other types of cancers [14,16].

Not only are these pesticides dangerous to the environment and humans, but they also cause the phenomenon of mirid resistance [17,18].

The search for alternatives to these synthetic chemical pesticides is therefore important. In this context, the use of local agro-resources is a serious avenue. Indeed, many plants have insecticidal properties that can be used in the control of cocoa mirids. Among these plants, neem (*Azadirachta indica*) is known and used for its insecticidal properties [19].

The scientific question guiding our study is this: What is the spectrum of insecticidal activity of neem oil? The objective of our work is therefore to contribute to a production of cocoa in Togo that respects the environment and human health. This will specifically assess the insecticidal and repellent properties of neem oil on fifth instar larvae and adults of *S. singularis*.

2. MATERIALS AND METHODS

2.1 Materials

2.1.1 Collection and rearing of *S. singularis*

The larvae were collected using fine brushes in boxes 11 cm of diameter on plots of certified organic cocoa that had not undergone any phytosanitary treatment. They are collected very early in the morning between 6 a.m. and 8 a.m. In the laboratory, the different larval stages were placed in different boxes measuring 45x30x30 cm. The upper end of the boxes was covered with a plastic net to facilitate ventilation. These larvae were used for the tests after an acclimatization period of 24 hours at ambient laboratory temperature ($t: 27\pm 2^{\circ}\text{C}$; RH: 80%). Adults are obtained by rearing part of the collected larvae. The larvae are placed in boxes measuring 45x30x30 cm and then fed with young leaves, branches, and cocoa pods until the emergence of adults. The inside of the boxes was lined with blotting paper. The homogeneous

batches of emerged adults were then used for the tests after an acclimatization period of 24 hours at laboratory temperature.

2.1.2 Neem oil

The neem oil was provided by BIOPHYTO, a Beninese company specializing in the research and exploitation of neem. The neem oil supplied is certified ORGANIC by ECOCERT, a certification organism.

2.2 Methods

2.2.1 Biological tests

2.2.1.1 Contact toxicity tests

The tests took place in vitro under laboratory conditions following a completely randomized experimental design with five (05) replications. Five (05) concentrations of neem oil were tested after presumptive tests to circumscribe the limit ranges of toxicity of the oil and Tween 80: 0.5; 1; 1.5; 2 and 2.5 $\mu\text{l.ml}^{-1}$. Neem oil was diluted in distilled water by adding Tween 80 as an emulsifier at a concentration of 0.1% non-toxic to insects. The solutions are prepared daily just before the tests.

A lot of twenty (20) insects was removed using a fine brush and then placed in a 45x30x30 cm box containing absorbent paper. The solution was then sprayed on the insects using a manual pressure sprayer of 15 ml capacity. After this operation, the insects were transferred to another box with a diameter of 11 cm containing pieces of young pods and young leaves placed on the absorbent paper lining the bottom of the box to serve as food for the treated insects. The upper end of the boxes was covered with a net. Each treatment, therefore, consists of an 11 cm diameter box containing twenty (20) insects. The absolute control consists of distilled water. The positive control consists of IMIDA 30 EC, a synthetic chemical insecticide composed of 30 g/l of imidacloprid, at a concentration of 3.3 $\mu\text{l.ml}^{-1}$. The mortality rate of the insects subjected to the various treatments was evaluated 24 h after the treatment.

Adjusted mortality was calculated using Abbott's formula [20]: $MC = (Me - Mt) / (100 - Mt) * 100$

Mc = Adjusted mortality; Me = sample mortality and Mt = untreated control mortality.

2.2.1.2 Repellency test

This test was carried out according to the preferential zone method on filter paper described by McDonald [21]. A disc of Wattman n°2 filter paper of 15 cm diameter was used. The filter paper was cut into two equal parts. The five concentrations of oil (0.5; 1; 1.5; 2 and 2.5 $\mu\text{l.ml}^{-1}$) were prepared by dilution in acetone. A volume of 0.5 ml of each solution was spread uniformly on one half of the disc and the other half received acetone only.

After fifteen minutes, the time necessary for the complete evaporation of acetone, the two halves of the discs were re-welded using adhesive tape. The filter paper disc thus reconstituted was placed in a 16 cm diameter box and a lot of twenty (20) insects was introduced into the center of the disc.

Five replications were performed for each concentration of the solution. After two hours, the number of insects present on the part of the filter paper treated with neem oil (Nt) and the number of those present on the part treated only with acetone (Nc) were recorded.

The percent repellency (PR) was then calculated using the following McDonald's formula [21]: $PR = (Nc - Nt) / (Nc + Nt) * 100$.

The PR helped to classify the solutions according to the repellent classes of McDonald et al. [21] from 0 to V.

2.2.2 Analysis of results

The results obtained were statistically analyzed using SPSS. 21 and the Student-Newman-Keuls test permitted discrimination between the means obtained.

The LC50s were calculated using the probit regression method.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Insecticidal effect of *A. indica* seed oil on 5th instar larvae and adults of *S. singularis*

Mortalities induced by neem seed oil on 5th-instar larvae and adults of *S. singularis* are presented in the Table 1.

The results show increasing mortality of all stages with increasing concentrations of neem seed oil. The concentration of 2.5 $\mu\text{l.ml}^{-1}$ caused the death of 100% of both 5th-instar larvae and adults. However, the concentration of 1 $\mu\text{l.ml}^{-1}$ caused the death of 54 \pm 3.98% of the 5th instar larvae whereas more than 1.5 $\mu\text{l.ml}^{-1}$ caused the death of 57.5 \pm 5.4% of adults (Table 1).

The LC₅₀ calculated using the probit regression method is given in the Table 2.

The results show that neem oil acts differently on the developmental stages of *S. singularis*. The calculated LC₅₀s increase according to the increasing stages of development of the insect. The LC₅₀s were 0.89 and 0.98 $\mu\text{l.ml}^{-1}$ respectively for 5th instar larvae and adults of *S. singularis* (Table 2)

3.1.2 Repellent effect of *A. indica* seed oil on 5th instar larvae and adults of *S. singularis*

Table 3 shows neem seed oil's repellency percentages (RP).

The results of our work show an increasing repellency percentage (RP) with increasing concentrations of neem seed oil for both 5th instar larvae and adults of *S. singularis*. The average RPs of neem seed oil on 5th instar larvae and adults are 23.6 and 28.8%, respectively (Table 3).

These results allowed us to classify neem oil according to McDonald's repellency classes which are presented in the Table 4.

The mean repellency percentages (MRP) calculated were 23.6 and 28.8 for 5th instar larvae and adults of *S. singularis*, respectively. These repellency percentages place neem seed oil in McDonald's Class II. Then, the neem seed oil is weakly repellent against both 5th instar larvae and adults of *S. singularis* (Table 4).

3.2 Discussion

Observed mortality increased with neem oil concentrations for both 5th instar larvae and adults of *S. singularis*. These results show that neem oil was toxic to both larvae and adults. However, calculated LC₅₀s revealed that neem oil is more toxic to larvae than adults. This toxicity is due to the seed oil's composition and mechanism of action. Indeed, the biological properties of neem are attributed to more than 135 active phytochemical constituents, including alkaloids, carotenoids, flavonoids, ketones, phenolics, steroids, and triterpenoids, which have been isolated from different parts of the neem plant. These compounds include azadirachtin, gedunin, isomargolonone, margolone, margolonone, nimbidine, nimbin, nimbolide, and salannine [22]. Among these compounds, azadirachtin which is the main active ingredient of neem oil is the cause of 90% of the effects on insects [23]. Azadirachtin and other compounds in neem products exhibit various modes of action against insects, such as antifeedant, growth inhibition, fertility suppression, disruption of oviposition, changes in morphology, and inhibition of development [24]. The contact toxicity obtained in our study can be explained by

Table 1. Average mortality of 5th instar larvae and adults of *S. singularis* after treatment with *A. indica* seed oil

Concentration ($\mu\text{l.ml}^{-1}$)	Mortality rate (%)	
	5 th instar larvae	Adults
0	0th	0f
0.5	39 \pm 4.54d	29 \pm 5.4e
1	54 \pm 3.98c	48 \pm 5.4d
1.5	67 \pm 5.05b	57.5 \pm 5.4b
2	78 \pm 6.21b	68.5 \pm 5.4b
2.5	100 \pm 00a	100 \pm 5.4a

Within the same column, the means assigned the same letter do not differ statistically from each other (Newman-Keuls test, $p=0.05$)

Table 2. CL₅₀ per contact neem seed oil tested on 5th instar larvae and adults of *S. singularis*

Stage	LC ₅₀ ($\mu\text{l.ml}^{-1}$)
5 th instar larvae	0.89
Adults	0.98

Table 3. Repellency Percentage of different concentrations of *A. indica* seed oil tested on 5th instar larvae and adults of *S. singularis*

Concentration ($\mu\text{l.ml}^{-1}$)	Repellency Percentage (RP)	
	5 th instar larvae	Adults
0.5	10	12
1	10	24
1.5	24	28
2	36	40
2.5	38	40
Mean PR	23.6	28.80

Table 4. Mean Repellency Percentage (MRP), McDonald classes, and properties of *A. indica* seed oil

Stage	MRP	McDonald's class	Properties
5 th instar larvae	23.6	II	Weakly repellent
Adults	28.8	II	Weakly repellent

the direct effect of azadirachtin on the cells and tissues of the insect. Indeed, azadirachtin is absorbed by cells and causes inhibition of cell division and protein synthesis. These effects manifest as mild muscle paralysis, midgut cell necrosis, loss of gut nidi (regenerative cells), and lack of midgut enzyme production [25]. Azadirachtin also affects insect respiration and digestion. This action has been demonstrated in *Anticarsia gemmatalis* [26]. Our results are similar to those of many studies conducted on the insecticidal effects of neem oil on mirid larvae and adults. Indeed, neem oil and the ethanolic extract of neem oil are effective against *S. singularis* and lead to mortality rates ranging from 72.5 to 97.5% by contact and 32.5 to 70.0% by ingestion [27]. Some studies demonstrated that aqueous extract of neem oil inhibit larval molting with an inhibition rate of more than 50% [28]. Moreover, this plant has an interest in the control of cocoa mirids because all its parts can be used because they contain azadirachtin.

Calculated average repellent rates revealed that neem oil has a weak repellent effect against both 5th instar larvae and adults of *S. singularis*. This repellent action is due to the effect of neem oil compounds on the olfactory sense organs of the insect. Although thousands of odorants potentially can interact with an insect's olfactory system, there are only four general mechanisms that have been shown so far to reduce an insect's contact with hosts: 1) the activation of olfactory receptors dedicated to aversion, 2) activation of pheromone receptors that cause aversion, 3) the inhibition of odorant receptors dedicated to attraction, and 4) prolonged

activation of odorant receptors participating in attraction [29]. Essential oil with oxygenated monoterpenes, sesquiterpene hydrocarbons bind insect "odor binding proteins" and make them confuse [30]. The repellent effect of neem oil is probably due to azadirachtin, a terpenoid [31, 32, 24]. Other studies have shown that neem oil has a repellent effect on several insects including *Tribolium castaneum* and *Bemisia tabacii* [32, 33]. The weak repellent effect of neem oil in our study could be explained by the concentration of the solutions used. Indeed, our results revealed that the rate of repellency is increasing with increasing concentrations of neem oil.

These results, therefore, show that neem oil can be an alternative to synthetic chemical pesticides in the fight against *S. singularis*.

4. CONCLUSION

The constraints of cocoa cultivation are numerous. Among these constraints, insect pressure, particularly *S. singularis*, causes the most loss. Chemical control of these insects leads to environmental pollution and endangers the lives of consumers. The search for alternatives to these synthetic chemical pesticides is essential. Thus, our work consisted in investigating the insecticidal and repellent properties of neem oil against 5th instar larvae and adults of *S. singularis*. Neem oil was insecticidal and weakly repellent against these stages of *S. singularis*. It can therefore be an alternative in the control of this insect. Our study will continue with field experiments to conclude on the effectiveness of neem oil.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Adden KA. Amélioration de la productivité des vergers de cacaoyers (*Theobroma cacao* Linn.) pour une gestion forestière durable au Togo. Ecole Supérieure d'Agronomie de l'Université de Lomé; 2017.
2. Koudjega T. TBK. Improvement of soils fertility management in cocoa plantations in Togo. In: Proceeding of the 7th International Symposium on Plant-soil Interactions at Low pH. Guangzhou, China. 2009;184-5.
3. Wessel M, Quist-Wessel PMF. Cocoa production in West Africa, a review and analysis of recent developments. NJAS - Wageningen J. Life Sci. 2015;74-75:1-7.
4. DSID Direction de la Statistique de l'Informatique et de la Documentation. Recensement des planteurs et des plantations de café cacao au Togo. Rapport finale. Lomé, Togo; 2014.
5. INSEED Institut National de la Statistique et des Etudes Economiques et de Développement. Comptes nationaux Togo 2010 – Comptes Finaux. Lomé, Togo; 2015.
6. Food and Agriculture Organisation of the United Nations. Online Database [Internet]; 2020. Available: <https://www.fao.org/faostat/en/#home>
7. Awudzi GK, Asamoah M, Owusu-Ansah F, Hadley P, Hatcher PE, Daymond AJ. Knowledge and perception of Ghanaian cocoa farmers on mirid control and their willingness to use forecasting systems. Int. J. Trop. Insect Sci. 2016; 36: 22-31.
8. Kumi E, Daymond AJ. Farmers' perceptions of the effectiveness of the Cocoa Disease and Pest Control Programme (CODAPEC) in Ghana and Its Effects on Poverty Reduction. Am J Exp Agric. 2015;7(5):257-74.
9. Adu-Acheampong R, Sarfo J, Appiah E, Nkansah A, Awudzi G, Obeng E, et al. Strategy for Insect Pest Control in Cocoa. Am J Exp Agric. 2015;6(6):416-23.
10. Khanna R, Gupta S. Agrochemicals as a potential cause of ground water pollution: A review. Int J Chem Stud. 2018;6(3):985-90.
11. Ntiamoah A, Afrane G. Environmental impacts of cocoa production and processing in Ghana: Life cycle assessment approach. J Clean Prod. 2008;16(16):1735-40.
12. Zainudin BH, Salleh S, Yaakob AS, Mohamed R. Comprehensive strategy for pesticide residue analysis in cocoa beans through qualitative and quantitative approach. Food Chem. 2022;368(July 2021):130778.
13. Okoffo ED, Fosu-Mensah BY, Gordon C. Contamination levels of organophosphorus and synthetic pyrethroid pesticides in cocoa beans from Ghana. Food Control. 2017;73:1371-8.
14. Polanco Rodríguez ÁG, Riba López MI, DelValls Casillas TÁ, Araujo León JA, Mahjoub O, Prusty AK. Monitoring of organochlorine pesticides in blood of women with uterine cervix cancer. Environ Pollut. 2017;220:853-62.
15. Attaullah M, Yousuf MJ, Shaikat S, Anjum SI, Ansari MJ, Buner ID, et al. Serum organochlorine pesticides residues and risk of cancer: A case-control study. Saudi J Biol Sci. 2018;25(7):1284-90.
16. Tabrez S, Priyadarshini M, Priyamvada S, Khan MS, NA A, Zaidi SK. Gene-environment interactions in heavy metal and pesticide carcinogenesis. Mutat Res - Genet Toxicol Environ Mutagen. 2014; 760:1-9.
17. Stanek J, Mairinger S, Wanek T, Kuntner C, Müller M, Langer O. Automated radiosynthesis of [18F]ciprofloxacin. Appl Radiat Isot. 2015;99:133-7.
18. Ninsin K, Adu-Acheampong R. The Ghana Cocoa Board (COCOBOD) approved insecticides, imidacloprid, thiamethoxam and bifenthrin, for the control of cocoa mirids (Hemiptera: Miridae): Implications for insecticide-resistance development in *Distantiella theobroma* (Dist.) and *Sahlbergella singularis*. Ghana J. Agric Sci. 2017;51:21-8.
19. Gnago J, Danho M, Agneroh T, Fofana I, Kohou A. Efficacité des extraits de neem (*Azadirachta indica*) et de papayer (*Carica papaya*) dans la lutte contre les insectes ravageurs du gombo (*Abelmoschus esculentus*) et du chou (*Brassica oleracea*) en Côte d'Ivoire. Int J Biol Chem Sci. 2011; 4(4).
20. WS A. A method of computing the effectiveness of an insecticide. Journal Econ Entomol. 1925;18:265-7.

21. McDonald LL, GRH, SR. Preliminary evaluation of new candidate materials as toxicants, repellents and attractants against stored product insects. *Agric. Res.* 882. Washington. 1970;183.
22. Ghosh V, Sugumar S, Mukherjee A, Chandrasekaran N. Neem (*Azadirachta indica*) Oils. *Essential Oils in Food Preservation, Flavor and Safety*. Elsevier Inc. 2015;593-599.
23. Campos EVR, de Oliveira JL, Pascoli M, de Lima R, Fraceto LF. Neem oil and crop protection: From now to the future. *Front Plant Sci.* 2016;7(October2016): 1-8.
24. Kumar R, Mehta S, Pathak SR. Bioactive constituents of neem. *Synthesis of Medicinal Agents from Plants*. Elsevier Ltd. 2018;75-103.
25. Mordue AJ, Nisbet AJ. Azadirachtin from the neem tree *Azadirachta indica*: Its action against insects. *An da Soc Entomológica do Bras.* 2000;29(4):615-32.
26. Farder-Gomes CF, Saravanan M, Martínez LC, Plata-Rueda A, Zanuncio JC, Serrão JE. Azadirachtin-based biopesticide affects the respiration and digestion in *Anticarsia gemmatalis* caterpillars. *Toxin Rev.* 2022; 41(2):466-75.
27. Mahob RJ, Taliedje DM, Mahot HC, Ngah IM, Enama SE, Cilas C, et al. Biocontrol of the brown cocoa mirids using neem oil and an ethanolic extract from neem under laboratory conditions. *African Entomol.* 2021;29(2):507-21.
28. Adu-Acheampong R. Evaluation of neem (*Azadirachta indica* A. Juss) seed extracts for the management of some cocoa mirid species. University of Ghana; 1997.
29. Clark JT, Ray A. Olfactory Mechanisms for Discovery of Odorants to Reduce Insect-Host Contact. *J Chem Ecol.* 2016; 42(9):919-30.
30. Mota TF, Silva CM d. A, Conceição M dos S, Fraga DBM, Brodskyn CI, Neto MF d. A, et al. Screening organic repellent compounds against *Lutzomyia longipalpis* (Diptera: Psychodidae) present in plant essential oils: Bioassay plus an in silico approach. *Acta Trop.* 2022; 229.
31. Borokar Archana, Pansare Tabassum DP. Review on Neem (*Azadirachta Indica* A. Juss .) As a Potential Source of Organic Pesticide. *Int J Ayurveda Pharma Res.* 2017;5(4):12-5.
32. Gul HT. Toxicological and repellent potential of some plant extracts against stored product insect pest, *Tribolium castaneum* (Herbst.) (Coleoptera: Tenebrionidae). *Int J Biosci.* 2013; 3(9):280-6.
33. Castillo-Sánchez LE, Jiménez-Osornio JJ, Delgado-Herrera MA, Candelaria-Martínez B, José J. Effects of the hexanic extract of neem *Azadirachta indica* against adult whitefly *Bemisia tabaci*. *J Entomol Zool Stud.* 2015;3(5):95-9.

© 2023 Ankou et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/101065>