



Effect of Potting Mixture and Polybag Size on Growth and Flowering of Asiatic Lily (*Lilium asiaticum* var. *Nashville*) under Protected Condition

Anupama Geethan ^{a++*}, Vijay Bahadur ^a
and Annjoe V Joseph ^a

^a Department of Horticulture, SHUATS, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i64692>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117325>

Original Research Article

Received: 16/03/2024
Accepted: 22/05/2024
Published: 24/05/2024

ABSTRACT

An experiment on recognizing the significance of cultivating lilies under polyhouse conditions, this study contributes the valuable insights to the cultivation practices of Asiatic lilies, focusing on the pivotal roles of potting mixture composition and polybag size. Healthy bulbs of each cultivar were freshly planted in various media compositions in different polybag size within a protected environment. From the present investigation, it is concluded that the treatment 12 x 12 polybag size combined with Garden soil + Vermicompost + Cocopeat (1:1:2) was found in superior among other treatments in terms of plant height (69.30cm), leaf length (8.60 cm), leaf width (3.03 cm), number of flower buds/plant (4.60), days to appearance of first flower bud (40.10), flower bud length (8.43

⁺⁺ M.Sc. Scholar;

*Corresponding author: E-mail: vijaybahadur@shuats.edu.in;

Cite as: Geethan, A., Bahadur, V., & Joseph, A. V. (2024). Effect of Potting Mixture and Polybag Size on Growth and Flowering of Asiatic Lily (*Lilium asiaticum* var. *Nashville*) under Protected Condition. *International Journal of Plant & Soil Science*, 36(6), 850–859. <https://doi.org/10.9734/ijpss/2024/v36i64692>

cm), stalk length (68.15 cm), weight of bulb per plant (64.75g), self life of flower spike (15.22 days), number of flowers per 250 m² (10032.75). The study concluded that, the lilies readily respond to various medium for cultivation and 12 x 12 Polybag combined with potting mixture Garden soil + Vermicompost + Cocopeat (1:1:2) observed to be the best for plant growth and flowering of Asiatic lily under protected condition.

Keywords: Cocopeat; FYM; liliium; media; Nashville; perlite; vermiculite.

1. INTRODUCTION

Lilium holds a significant position as a commercial bulbous cut flower cultivated worldwide. Belonging to the family *Liliaceae*, the genus *Lilium* encompasses over 80 species spread across North America, Asia, and Europe, showcasing notable diversity in plant structure, flower characteristics, bulb types, and fragrances [1,2]. Lilium, ranks fourth among the bulbous plants after Tulipa, Gladiolus, and Narcissus [3]. Its popularity stems from its exquisite blooms, diverse varieties, and versatility in landscaping, cut flower production, and ornamental gardening [4]. With its wide array of colours, shapes, and sizes, Lilium continues to be cherished by growers, florists, and enthusiasts alike, contributing to the vibrant and diverse floral industry [5].

Lilium, renowned for its distinctive bulbous flowers, stands as a highly sought-after cut flower in the global market. So, the commercial cultivation of Asiatic hybrid lilies is expanding steadily in India. Its popularity stems from its diverse colour palette, robust shelf life, and the ease with which it thrives under both open field and shaded conditions. Noteworthy is its ability to rehydrate effectively post long-distance transportation when provided with suitable cut flower food, thus ensuring its freshness for consumers [6]. Soil-borne diseases and pests pose significant challenges in the production of cut flower Lilium, often leading to a notable decline in flower quality. As a result, soilless cultivation has emerged as the preferred method for ornamental plants among modern growing techniques [7].

Cocopeat, a byproduct of coconut processing, is a fibrous material known for its excellent moisture retention and soil aeration properties [8]. Vermiculite, a mineral that expands when heated, is valued for its ability to improve soil structure and moisture retention, making it popular in gardening and horticulture [9]. Perlite, another lightweight mineral, is prized for its exceptional drainage properties and aeration

benefits, commonly used in potting mixes to promote healthy root growth [10]. Vermicompost, on the other hand, is compost produced through the digestion of organic waste by earthworms. Rich in nutrients and beneficial microorganisms, vermicompost enhances soil fertility, improves soil structure, and promotes vigorous plant growth [2]. Each of these soil amendments plays a unique role in improving soil health and supporting plant growth in various gardening and agricultural applications.

The characteristics of the growing media utilized in soilless cultivation directly or indirectly impact both yield and quality and lilies readily respond to various medium for cultivation [11]. Thus, recognizing the significance of cultivating lilies under polyhouse conditions, this study contributes the valuable insights to the cultivation practices of Asiatic lilies, focusing on the pivotal roles of potting mixture composition and polybag size. By elucidating the optimal conditions for growth and flowering, this research aims to empower growers in Prayagraj and similar agro-climatic regions to harness the full potential of *Lilium asiaticum* as a prominent cut flower in the international trade market.

2. MATERIALS AND METHODS

The present investigation was carried out in the form of field experiment in Horticultural Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh, India during 2023-2024 under protected condition. The experiment was laid out in Completely Randomized Design with twelve treatments and three replications. The experiment includes the combination of various potting mixtures viz., garden soil, cocopeat, vermicompost, FYM, vermiculite and perlite in varying proportions and different polybag sizes (6, 31 & 35). The metrological observations during the experimental period of November 2023 to March 2024 is represented in Table 1.

Different media composition used for filling of pots were; T₁: Garden soil+ FYM+Cocopeat (1:1:2) in 8x8 polybag size, T₂: Garden soil+ Vermicompost +Cocopeat (1:1:2) in 8x8 polybag size, T₃: Garden soil+ Cocopeat + Vermiculite (1:2:1) in 8x8 polybag size, T₄: Garden soil+ Cocopeat + Vermiculite + Perlite (1:2:1:1) in 8x8 polybag size, T₅: Garden soil+ FYM+Cocopeat (1:1:2) in 10x10 polybag size, T₆: Garden soil+ Vermicompost +Cocopeat (1:1:2) in 10x10 polybag size, T₇: Garden soil+ Cocopeat + Vermiculite (1:2:1) in 10x10 polybag size, T₈: Garden soil+ Cocopeat + Vermiculite + Perlite (1:2:1:1) in 10x10 polybag size, T₉: Garden soil+ FYM+Cocopeat (1:1:2) in 12x12 polybag size, T₁₀: Garden soil+ Vermicompost +Cocopeat (1:1:2) in 12x12 polybag size, T₁₁: Garden soil+ Cocopeat + Vermiculite (1:2:1) in 12x12 polybag size and T₁₂: Garden soil+ Cocopeat + Vermiculite + Perlite (1:2:1:1) in 12x12 polybag size. Observations were recorded for the six plants within each treatment at 20-days interval. Statistical analysis was conducted using Fisher's analysis of variance (ANOVA) technique, in accordance with the guidelines outlined by Gomez and Gomez [12].

3. RESULTS AND DISCUSSION

A. Impact of Potting Mixture & Polybag Size on Vegetative Parameters

Vegetative parameters are essential for assessing the growth and development of plants, understanding their response to different environmental conditions, and evaluating the effectiveness of various treatments and the results on the effect of different growing media and polybag sizes of these vegetative parameters on Asiatic lily are represented in Table 2.

Plant height was observed maximum (69.30 cm) in T₁₀ - 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) followed by T₁₁ - 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (66.27 cm) whereas, minimum (60.73 cm) was reported in T₂ - 8 x 8 - Garden soil + Vermicompost + Cocopeat (1:1:2). The significant increase in plant height observed in the treatment using a potting mixture of Garden soil, Vermicompost, and Cocopeat (1:1:2) in 12 x 12 polybags can be attributed to the unique mineral composition of the substrates. Cocopeat, rich in essential minerals for crop plant growth, likely played a pivotal role in this enhancement [13] demonstrated that a blend comprising 30%

vermicompost and 70% cocopeat resulted in the tallest plants (81.7 cm) in liliium, while substrates without cocopeat led to shorter heights (67.5 cm).

The Number of leaves per plant (54.17) was observed maximum in T₁₁- 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) followed by T₁₀- 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) (54.08) whereas, minimum (47.63) was reported in T₃- 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). Vermicompost, known for its water retention and nutrient uptake, combined with cocopeat, boosts organic nutrients for liliium growth [14] found more leaves with vermicompost in 'Navona' lilies, while [15] noted highest leaf counts in a mix of rice hull, sawdust, and vermicompost.

The Leaf length (8.60 cm) was observed maximum in T₁₀- 12 x 12- Garden soil+ Vermicompost + Cocopeat (1:1:2) followed by T₁₂- 12 x 12- Garden soil+ Cocopeat + Vermiculite + Perlite (1:2:1:1) (8.17 cm) whereas, minimum (7.03 cm) was reported in T₃- 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). Photosynthesis is crucial for plant growth, as it produces their food. More leaves usually mean increased photosynthesis, driving overall growth. Minerals absorbed by plants enhance leaf size, boosting photosynthesis and yields. This aligns with Choi et al. [15] in liliium [16] found cocopeat-amended mediums led to more leaves, area, and better yields in liliium.

The maximum Leaf Width (3.03 cm) was observed with T₁₀- 12 x 12- Garden soil+ Vermicompost +Cocopeat (1:1:2) which was at par with T₁₁- 12 x 12 -Garden soil+ Cocopeat + Vermiculite (1:2:1) (2.97 cm) whereas, minimum (2.47 cm) was reported in T₁- 8 x 8 - Garden soil+ FYM + Cocopeat (1:1:2). Potassium in vermicompost boosts plant metabolism, aiding in amino acid and protein production crucial for growth. It also supports photosynthetic pigment emergence, enlarging leaf surfaces [14] found similar results in liliium, indicating that vermicompost and coirpith enhance leaf characteristics, signaling robust growth and higher yields.

B. Impact of Potting Mixture & Polybag Size on Floral Parameters

Floral parameters provides the ornamental value, and overall health of plants, particularly in the economic view of liliium plants and the results on

the effect of different growing media and polybag sizes on Asiatic lily are represented in Table 3.

The maximum Number of Flower Buds/Plant (4.60) was observed with T₁₀- 12 x 12- Garden soil+ Vermicompost +Cocopeat (1:1:2) which was at par with T₁₁- 12 x 12 -Garden soil+ Cocopeat + Vermiculite (1:2:1) (3.93) whereas, minimum (2.23) was reported in T₃- 8 x 8 - Garden soil+ Cocopeat + Vermiculite (1:2:1). Optimal conditions from vermicompost and cocopeat enhance plant health and flower production. Vermicompost, a rich nutrient source, boosts growth, while cocopeat provides ideal moisture and aeration for early and improved flowering. This fosters a higher flower count per plant, as seen in studies like [14] on Asiatic hybrid lily 'Navona'.

The minimum Days to appearance of flower bud (40.10) was observed with T₁₀- 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) which was at par with T₁₁- 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (41.20) whereas, maximum (50.83) was reported in T₃ - 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). Variations in flowering initiation time may be due to cocopeat and vermicompost benefits. Cocopeat offers aeration and moisture, promoting early flowering, while vermicompost supplies vital nutrients. This aligns with [9] findings. Similar results were reported by Bhatia et al. [17], noting shorter flower bud formation time (94.41 days) in carnations with soil, vermicompost, and cocopeat (in 2:1:1 ratio) [18] also observed quicker flower bud formation in anthuriums with cocopeat and vermicompost.

The maximum Flower bud length (8.43 cm) was observed with T₁₀ - 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) which was at par with T₁₁ - 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (7.99 cm) whereas, minimum (6.64 cm) was reported in T₂- 8 x 8 - Garden soil + Vermicompost + Cocopeat. Flower quality significantly impacts the appeal of cut flowers, determined by factors like floret arrangement, size, and number on the spike. Research by Rani et al. [19] on LA Liliu cultivars found that mediums with cocopeat and vermicompost led to early bud visibility, longer stalks, and flower bud lengths compared to natural soil. Similar results were noted by Rajvanshi et al. [20] and [21], suggesting gibberellins in vermicompost may regulate the reproductive phase, enhancing stalk and bud lengths. Vermicompost, rich in micronutrients

and macronutrients, promotes plant growth, improving flower characteristics [22] experiments on LA hybrid lilies support this, indicating cocopeat and vermicompost amendments enhance plant growth, resulting in longer flower bud and stalk lengths, and prolonged vase life.

The maximum Stalk Length (68.15 cm) was observed with T₁₀ - 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) which was at par with T₁₁- 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (67.30 cm) whereas, minimum (60.10 cm) was reported in T₃ - 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). Stalk length is crucial for export quality, as per [23], cocopeat's carbon content and porosity foster beneficial microorganism growth, enhancing nitrogen fixation, phosphorus solubilization, and phytohormone production for better plant growth [9] found vermicompost and cocofibre combo maximized anthurium stalk length due to improved aeration and moisture, a conclusion supported by Bhatia et al. [17,24,25].

The maximum Self-life of flower spike (15.22 days) was observed with T₁₀ - 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) which was at par with T₁₁ - 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (14.58 days) whereas, minimum (10.02 days) was reported in T₃ - 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). The prolonged shelf life of cut flowers is linked to stored carbohydrate content, crucial for longevity. Cocopeat and vermicompost in the growing medium help maintain high water content and nutrient uptake, enhancing flower durability [24]. Prisa et al. [26] observed improved flower quality and longevity in Asiatic hybrids on new substrates compared to traditional ones [17] found the longest flowering duration (70.56 days) in soil, vermicompost, and cocopeat mediums, consistent with Jawaharlal et al. [18] report of 87.50 days for anthuriums in cocopeat.

C. Impact of Potting Mixture & Polybag Size on Bulb and Yield Parameters

Bulb and yield parameters encompass on various aspects such as the weight, diameter and yield per plant, indicating the quantity of harvested produce obtained from individual plants, and the total yield, which represents the overall quantity of harvested produce from the entire cultivation area. The results on the effect of different growing media and polybag size on Asiatic lily are represented in Table 3.

The maximum bulb diameter (6.12 cm) was observed with T₁₀- 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) (64.75) which was at par with T₁₁ - 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (5.78 cm) whereas, minimum (3.98 cm) was reported in T₃ - 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). Differences in liliium bulb diameter may be due to higher organic matter percentages in vermicompost and cocopeat media, enhancing nutrient availability and promoting cell division and size elongation [27] noted larger flower diameters in peat and vermicompost mixes, improving media physicochemical characteristics. Similar findings were reported by [10,28,29,17,30,20].

The maximum weight of bulb per plant (g) was observed with T₁₀- 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) (64.75) which was at par with T₁₁ - 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (62.18) whereas, minimum was reported in T₃ - 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1) (43.91). The rise in bulb weight may be attributed to Gibberellin in vermicompost, promoting cell

division, elongation, and differentiation, along with enhancing protein and nucleic acid uptake, resulting in increased bulb weight. Similar outcomes were observed by [31,32,33]. Nikrazm et al. [16] noted increased bulb size, measured by perimeter, in cocopeat mediums for liliium [34] found maximum bulb size in soil with cocopeat and vermicompost (2:1:1, v/v) for liliium Asiatic hybrids.

The maximum Weight of bulb per plant (0.064 kg) was observed with T₁₀- 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) followed by T₁₁ - 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (0.062 kg) whereas, minimum (0.043 kg) was reported in T₃ - 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). Vermicompost-amended mediums in large polybags enhance vegetative and flower parameters, likely leading to heavier bulbs. Gibberellin from vermicompost enhances cell division and elongation, potentially increasing protein and nucleic acid uptake. Similar results were reported for Oriental lilies with cocopeat-containing media [35].

Table 1. Metrological observations during the experimental period of Nov. 2023 - Mar. 2024

Month and year	Week	Temperature (°C)		Relative humidity (%)	
		Max.	Min.	Max.	Min.
November, 2023	1 st week	36.35	22.89	89.84	49.70
	2 nd week	27.90	25.39	86.27	47.70
	3 rd week	27.30	13.04	85.49	39.13
	4 th week	25.22	29.99	92.99	61.99
December, 2023	1 st week	33.20	13.84	86.56	47.70
	2 nd week	31.75	10.56	89.41	53.70
	3 rd week	27.09	10.99	92.56	60.41
	4 th week	21.00	9.59	97.70	64.56
January, 2024	1 st week	24.77	15.24	94.70	58.99
	2 nd week	21.32	10.70	94.56	74.70
	3 rd week	18.92	9.84	98.27	68.13
	4 th week	22.00	14.84	95.27	75.27
February, 2024	1 st week	27.18	12.88	82.13	47.13
	2 nd week	29.78	14.99	77.56	47.28
	3 rd week	31.21	13.85	80.42	46.56
	4 th week	35.86	15.70	77.99	44.70
March, 2024	1 st week	37.66	15.62	77.56	42.42
	2 nd week	37.92	16.88	78.85	42.70
	3 rd week	35.61	16.42	82.13	54.28
	4 th week	34.38	20.22	70.56	49.59

Table 2. Effect of potting mixture and polybag size on vegetative growth and floral parameters of Asiatic lily

Tr.	Plant height (cm)	Number of leaves/plant	Leaf length (cm)	Leaf width (cm)	Number of Flower Buds/Plant	Days to appearance of flower bud	Flower bud length (cm)	Stalk Length (cm)	Self-life of flower (days)
T ₁	64.53	51.53	7.40	2.47	2.55	50.25	7.33	62.30	10.33
T ₂	60.73	51.20	8.07	2.60	2.73	46.50	6.64	61.45	11.83
T ₃	61.40	47.63	7.03	2.73	2.23	50.83	6.73	60.10	10.02
T ₄	61.27	50.74	7.97	2.80	3.87	49.53	7.20	61.25	12.12
T ₅	64.60	52.88	7.40	2.77	3.03	44.12	7.37	63.11	10.49
T ₆	62.73	51.27	6.83	2.57	3.27	48.73	7.67	61.80	11.62
T ₇	63.40	52.43	7.83	2.87	3.08	45.97	7.53	63.30	12.18
T ₈	64.63	51.93	8.10	2.70	3.15	50.20	7.43	63.28	10.55
T ₉	65.73	53.97	7.90	2.63	3.21	44.40	7.83	64.50	13.77
T ₁₀	69.30	54.18	8.60	3.03	4.60	40.10	8.43	68.15	15.22
T ₁₁	66.27	54.07	7.97	2.97	3.93	41.20	7.99	67.30	14.58
T ₁₂	61.33	54.00	8.17	2.67	3.37	42.50	7.64	66.12	13.26
F-Test	S	S	S	S	S	S	S	S	S
S.Ed (±)	1.97	2.62	0.46	0.19	0.19	0.99	0.35	1.50	0.81
CD _(0.05)	4.09	5.45	0.97	0.52	0.41	2.05	0.73	3.13	3.24

Table 3. Effect of potting mixture and polybag size on bulb and yield parameters of Asiatic lily

Treatments	Diameter of the Bulb (cm)	Weight of bulb per plant (g)	Weight of bulb per plant (kg)	Weight of bulb per 250 m ² (kg)	Number of Flowers per 250 m ²
T ₁	4.20	49.88	0.049	104.54	7338.78
T ₂	4.52	47.02	0.047	106.69	8528.02
T ₃	3.98	43.91	0.043	102.37	5312.34
T ₄	4.11	52.27	0.052	118.61	7489.22
T ₅	4.29	56.38	0.056	124.14	8094.38
T ₆	4.75	58.33	0.058	127.25	9442.86
T ₇	4.52	56.92	0.056	119.32	9298.04
T ₈	4.47	54.07	0.054	116.29	9342.18
T ₉	5.21	58.61	0.058	130.55	9574.91
T ₁₀	6.12	64.75	0.064	143.81	10032.75
T ₁₁	5.78	62.18	0.062	140.38	9867.12
T ₁₂	5.45	59.55	0.059	138.24	9612.88
F-Test	S	S	S	S	S
S.Ed (±)	0.11	1.98	1.56	6.05	543.95
CD (5%)	0.24	4.22	4.01	14.23	1091.12

The weight of bulbs per plant per 250m² (143.81 kg) was observed with T₁₀- 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) followed by T₁₁ - 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (140.38 kg) whereas, minimum (102.37 kg) was reported in T₃ - 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). Maximum bulblet weight was achieved in soil amended with garden soil, cocopeat, and vermicompost in 12 x 12 polybags due to enhanced carbohydrate translocation underground, facilitating bulblet formation. This nutrient-rich combination promotes stem elongation via increased cell division. Soil microflora continuously mineralize complex substances, synthesizing biologically active metabolites, potentially leading to more leaves per plant and heavier bulb weight [24].

The maximum number of flowers per 250 m² (10032.75) was observed with T₁₀- 12 x 12 - Garden soil + Vermicompost + Cocopeat (1:1:2) followed by T₁₁ - 12 x 12 - Garden soil + Cocopeat + Vermiculite (1:2:1) (9867.12 kg) whereas, minimum (5312.34) was reported in T₃ - 8 x 8 - Garden soil + Cocopeat + Vermiculite (1:2:1). Cocopeat and vermicompost positively impact flower numbers due to their enhanced structure, nutrient content, and moisture retention, especially in larger polybags [36] found the highest net returns in gerbera flowers with a cocopeat and vermicompost blend (1:1 ratio), consistent with Sekar and Sujata [25] for gerberas. Similar outcomes were seen by Krause [37] and [38] in *Lilium longiflorum*, [39] in Asiatic cultivars, and [39] in Oriental lilies [40-43].

4. CONCLUSION

The present study indicates that the treatment involving garden soil, cocopeat, and vermicompost in 12 x 12 polybags exhibited superior growth and yield attributes compared to other treatments. This superiority is likely due to their positive effects on soil structure, nutrient availability, and moisture retention, which significantly influence both vegetative and reproductive characteristics of liliium plants. The utilization of cocopeat and vermicompost as amendments has been demonstrated to have a beneficial impact on various aspects of plant growth, such as plant height, leaf number and dimensions, flower characteristics, bulb weight, and shelf life. By optimizing the composition of the medium, growers have the potential to enhance crop productivity, quality, and marketability, meeting the requirements of both

local and export markets. Furthermore, insights gained from this study contribute to our understanding of the role of organic amendments in sustainable horticultural practices and emphasize their potential for improving crop production and post-harvest traits.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Grassotti A, Gimelli F. Bulb and cut flower production in the genus *Lilium*: Current status and the future. *Acta Horticulturae*. 2011;900:21–36.
2. De Jong PC. Some notes on the evolution of lilies. *Lily yearbook*, North American Lily Society. 1974;27:23–8.
3. De Hertogh AA. Marketing and research requirements for liliium in North America. *Acta Horticulturae*. 1996;414:17-24.
4. Faust JE, Dole JM. Major cut flowers. *Cut Flowers and Foliages*; CAB International: Boston, MA, USA. 2021;48-149.
5. Boddy K. *Blooming Flowers: A Seasonal History of Plants and People*. Yale University Press; 2020.
6. Mishra V, Dwivedi SK. Postharvest management of fresh cut flowers. *Postharvest Biology and Technology of Horticultural Crops. Principles and Practices for Quality Maintenance*. Apple Academic Press and CRC Press Taylor and Francis Group. Waretown, NJ, USA. 2015;347-399.
7. Karaguzel O. Effects of different growing media on the cut flower performances of oriental two *Lilium* varieties. *International Journal of Agricultural and Biological Engineering*. 2020;13(5):85-92.
8. Udayana SK, Naorem A, Singh NA. The multipurpose utilization of coconut by-products in agriculture: prospects and concerns. *Int J Curr Microbiol Appl Sci*. 2017;6:1408-1415.
9. Kalirana K, Dubey P. Effect of substrates on anthurium culture. *The Asian Journal of Horticulture*. 2008;(1):165-166.
10. Ahmed M, Khan MF, Hamid A. Effect of urea, DAP and FYM on growth and flowering of dahlia (*Dahlia variabilis*). *International Journal of Agriculture and Biology*. 2004;6(2):393-395.

11. Samantaray P, Nayak A, Pradhan I, Pattanaik A, Sahoo R, Mohanty S. Liliium: A High-Value Cut Flower Production Guide for Lucrative Return. *Journal of Scientific Research and Reports*. 2024;30(6):67-86.
12. Gomez KA, Gomez AA. *Statistical Procedures for Agriculture Research*, John Wiley and Sons. 1984;664–5.
13. Seyedi M, Allahyari MS. The impact of perlite and cocopeat as the growth media on liliium. *Asian Journal of Experimental Biology and Sciences*. 2012;3(3):502-505.
14. Moghadam ARL, Ardebili ZO, Saidi F. Vermicompost induced changes in growth and development of Liliium Asiatic hybrid var. Navona. *African Journal of Agricultural Research*. 2012;7(17):2609-2621.
15. Choi JJ, Lee JS, Choi JM. Effect of physicochemical properties of growing media on growth, nutrient uptake and soil nutrient concentration in pot plant production of asiatic hybrid lily 'Orange Pixie'. *Journal of the Korean Society for Horticultural Science*. 2002;43(6):747-753.
16. Nikrazm R, Ajirlou SA, Khaligy A, Tabatabaei SJ. Effect of different media on vegetative growth of two Liliium ciltivars in soilless culture. *Journal of Science and Technology of Greenhouse Culture*. 2011; 2(6):1-9.
17. Bhatia S, Gupta YC, Dhiman SR. Effect of growing media and fertilizers on growth and flowering of carnation under protected condition. *Journal of Ornamental Horticulture*. 2004;7(2): 174-178.
18. Jawaharlal M, Joshua JP, Arumugam T, Vijaykumar M. Standardization of growing media for anthurium (*Anthurium andreanum*) cv. 'Temptation' under shade net house. *South Indian Horticulture*. 2001; 49(7):323-325.
19. Rani N, Kumar R, Dhatt KK. Effect of nitrogen levels and growing media on growth, flowering and bulb production of Liliium cultivars. *Journal of Oranamental Horticulture*. 2005;8(1):36-40.
20. Rajvanshi SK, Dwivedi DH, Meena ML, Ram RB. Effect of various organic potting mixtures on vegetative growth and flowering of cosmos (*Cosmos sulphureas* Linn.) and gaillardia (*Gaillardia pulchella* Linn.) *Plant Archives*. 2011;11(1):127-130.
21. Bohra M, Kumar A. Studies on effect of organic manures and bioinoculants on vegetative and floral attributes of chrysanthemum cv. Little Darling. *International Journal of Life Sciences*. 2014;9(3):1007-1010.
22. Mohammadi S, Allahyari MS. The impact of perlite and cocopeat as the growth media on liliium. *Asian Journal of Experimental Biology and Sciences*. 2012; 3(3):502-505.
23. Noguera P, Abad M, Noguera V. Coconut coir waste: a new and viable ecologically, friendly peat substitute. *Acta Horticulture*. 2000;517:279 283.
24. Chauhan RV, Varu DK, Kava KP, Savaliya VM. Effect of different media on growth, flowering and cut flower yield of gerbera under protected condition. *The Asian Journal of Horticulture*. 2014;9(1):228 231.
25. Sekar K, Sujata A. Effect of growing media and GA₃ on growth and flowering of gerbera (*Gerbera jamesonii* H. Bolus.) under naturally ventilated greenhouse. *South Indian Horticulture*. 2001;49:338-339.
26. Prisa D, Burchi G, Antonetti M, Teani A. Use of organic or inorganic substrates for reducing the use of peat and improving the quality of bulbs inflorescences in asiatic lily. *Acta Horticulturae*. 2011;900:143-148.
27. Fakhri M, Maloupa E, Gerasopoulos D. Effect of substrate and frequency of irrigation in yield and quality of three *Gerbera jamesonii* cultivars, *Acta Horticulture*. 1995;40(8):41-45.
28. Anuje AA, Dalal SR, Gunge VS, Deshpande RM. Effect of growing media on growth, flowering and yield of gerbera under polyhouse conditions. *Orissa Journal of Horticulture*. 2004;32(2): 106-108.
29. Barman Bharathi TU, Medhi RP. Effect of media and nutrition on growth and flowering of *Cymbidium* hybrid 'HC. Arora'. *Indian Journal of Horticulture*. 2012;69 (3):395-398.
30. Khalaj MA, Amiri M, Sindhu SS, Awani Kumar Singh and Singh JP. Response of different growing media on the growth and yield of gerbera (*Gerbera jamesonii*) under hydroponic open system. *The Asian Journal of Horticulture*. 2011;6(2):484-487.
31. Jha S, Sharma GL, Dikshit SN, Patel KL, Tirkey T, Sarnaik DA. Effect of vermicompost and FYM in combination with inorganic fertilizer on growth, yield and flower quality of gladiolus (*Gladiolus hybridus*). *Journal of Soils and Crops*. 2012;22(2):253-257.

32. Lal S, Lakhawat SS, Choudary MK. Effect of organic manures and biofertilizers on growth, flowering and bulb production in tuberose cv. Single. Indian Journal of Horticulture. 2010;67(4): 554-556.
33. Shankar D, Dubey P. Effect of NPK, FYM and NPK + FYM on growth, flowering and corm yield of gladiolus when propagated through cormels. Journal of Soils and Crops. 2005;15(1):34-38.
34. Klasman R, Moreira D and Benedetto A. Cultivation of Asiatic hybrid of Lilium sp. in three different substrates. Revistala Facultad de Agronomia Universidad de Buenos Aires. 2002; 22(1):79-83.
35. Treder J. Growth and quality of oriental lilies of different fertilization levels. Acta Horticulturae. 2005;6673 297-302.
36. Barreto MS, Jagtap KB. Assessment of substrates for economical production of gerbera (*Gerbera jamesonii* Bolus ex Hooker F.) flowers under protected cultivation. Journal of Ornamental Horticulture. 2006;9(2):136-138.
37. Krause J. Reproductive ability of some *Lilium longiflorum* varieties propagated from bulb scales. Prace-z –Zakresu- Nauk-Rolniczych. 1996;81:267-273.
38. Matsuo E, Arisumi K. Varietal differences in the development of the scale bulblet in the easter lily. Memories of the Faculty of Agriculture, Kagoshima University. 1979; 15:79-85.
39. Singh A. Effect of position, plant growth regulators and growing media on bulblet production of liliium hybrids. Ph.D. Thesis, Dr. Y. S. Parmar University of Horticulture and Forestry, Nauni, Solan (H.P), India; 2002.
40. Ali U, Sajid N, Khalid A, Riaz L, Rabbani MM, Syed JH, Malik RN. A review on vermicomposting of organic wastes. Environmental Progress & Sustainable Energy. 2015;34(4):1050-1062.
41. Caron J, Price JS, Rochefort L. Physical properties of organic soil: adapting mineral soil concepts to horticultural growing media and histosol characterization. Vadose Zone Journal. 2015;14(6):1-14.
42. Kazemi F, Mohorko R. Review on the roles and effects of growing media on plant performance in green roofs in world climates. Urban Forestry & Urban Greening. 2017;23:13-26.
43. Sharma P, Sharma YD, Dhiman SR, Gupta YC. Effect of growth regulators and growing media on propagation of oriental lily hybrids through scaling. Journal of Ornamental Horticulture. 2007; 10(3):148-152.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/117325>