



Performance of Different Varieties of Gladiolus under Prayagraj agro-Climatic conditions

G. Nikhil Kumar ^{a++*}, Samir E. Topno ^{a#}
and Vijay Bahadur ^{at}

^a Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj – 211007, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i7987>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118054>

Original Research Article

Received: 02/04/2024

Accepted: 06/06/2024

Published: 10/06/2024

ABSTRACT

An experiment was carried out in the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, from November, 2023 to March, 2024. The experiment was conducted in Randomized Block Design (RBD) with fifteen varieties, replicated thrice. It is concluded that among the different varieties of gladiolus variety V₆ (Pusa Suhagin) performed significantly better in plant height (90.69 cm), number of leaves (10), leaf area index (0.22), spike length (113.6 cm), more number of spikes per plant (2.83), more number of florets per spike (16), however less number of days taken for first sprouting recorded in V₁₅ (Snow Board, 7 days) and Vase life highest in variety V₉ (Arka Pratham, 11.66 days).

⁺⁺ M.Sc. Scholar;

[#] Assistant Professor;

[†] Associate Professor & Head;

^{*}Corresponding author: Email: nikhilkumarlucky1999@gmail.com;

Cite as: Kumar, G. Nikhil, Samir E. Topno, and Vijay Bahadur. 2024. "Performance of Different Varieties of Gladiolus under Prayagraj Agro-Climatic Conditions". *Journal of Advances in Biology & Biotechnology* 27 (7):270-78. <https://doi.org/10.9734/jabb/2024/v27i7987>.

Keywords: *Variety gladiolus; Pusa Suhagin; Arka Pratham; flowering; growth.*

1. INTRODUCTION

“Flowers are one of the most beautiful gifts that nature has given to man. Their delightful smell makes people happy and joyful. Flowers are used for a number of additional purposes, including garlands, bouquets and interior decor in an old and religious culture. In India, where they are deeply associated with nearly all social occasions and celebrations” [1,2,3]. In addition, cultivating attractive plants and flowers is crucial for sustaining the ecological balance and reducing pollution in the environment.

“Gladiolus is a flower of glamour and perfection which is known as the queen of bulbous flowers due to its flower spikes with florets of massive form, brilliant colors, attractive shapes, varying size and excellent shelf life. The modern varieties are botanically known as *Gladiolus grandiflorus* belonging to the family Iridaceae. Gladiolus is grown as flower bed in gardens and used in floral arrangements for interior decoration as well as making high quality bouquets” [4].

“Gladiolus is rated fourth in worldwide trade, third in India's cut flower output and sixth in loose flower production” [5]. “Gladiolus flower cultivation generates six times the profit of rice production. Gladiolus cut flowers are produced by different countries such as USA, Holland, Italy, France, Poland, Bulgaria, Brazil, Australia and also Israel [6-12]. Apart from this in Europe, for over 100 years Gladiolus has been popular, whereas it has been introduced recently in India. India has suitable agro-climatic conditions for gladiolus cultivation; it is commercially cultivated in Himachal Pradesh, Sikkim, Karnataka, Uttar Pradesh, West Bengal, Tamil Nadu, Punjab and Delhi. In the eastern states like Tripura, Assam, Manipur, Meghalaya and Nagaland, this flower has established itself as a commercial proposition. There is a sizeable area under gladiolus in Jammu-Kashmir, Andhra Pradesh and Gujarat also [13-16].

During the last two decades the commercial production of gladiolus spikes has become a very popular flowering plant in India. The cultivation of gladiolus has the potential to change the economic scenario of farmers especially under Northern Indian plains, Uttar Pradesh, Tamil Nadu and Pondicherry. It is also necessary to examine the efficiency of different varieties and varieties available in the market [17-19]. So, all

these cultivars need to be tested for their performance, colors and color combinations, floral characteristics like spike length, more number and better size of floret, increase vase life etc [20-23].

Considering the present status, it is necessary to improve both quantitative and qualitative features in order to capitalize on internal and exotic demand. Climate and soil conditions have a significant impact on plant nutrition, development and subsequent survival while growing the gladiolus crop. Different climatic conditions have a significant impact on the growth and yield performance of Gladiolus. Commercial cultivation of gladiolus is becoming more popular, with large-scale production practiced mostly during the winter season. This experiment was conducted to evaluate the performance of different varieties of gladiolus.

2. MATERIALS AND METHODS

The field experiment entitled Performance of different varieties of gladiolus under Prayagraj agroclimatic conditions was carried out at the Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Naini, Prayagraj, during November, 2023 to March, 2024. The experimental field is situated on the left side of Prayagraj - Rewa Road, near the river Yamuna, and approximately 7 km from Prayagraj city.

2.1 Geographical Location and Climate Condition

Geographically, Prayagraj is situated in the south-eastern part of Uttar Pradesh. It lies between 25° 47' North latitudes and 82° 21' East longitudes. The area of Prayagraj district comes under agro-climatic zone V (Upper Gangetic Plain region) and sub-zone of Central Plains. The climate ranges from dry sub-humid to semi-arid and the soil is alluvium calcareous sandy loam. The district experiences an average maximum temperature range between 43° - 47°C which may go as high as 48°C during peak summers (May-June). The minimum average temperature is 2-4°C, which may fall as low as 1°C during peak winter months (December-January). The average rainfall of the district is 960 mm and the monsoon season is spread between July-September.

2.2 Experimental Details

The trail was laid out in a Randomized block design with fifteen varieties replicated thrice. Spacing 1 m x 1m. Fifteen different varieties used Priscilla, Joshika, Dhanvantari, Phule Neel Rekha, Souvik Biscuit, Pusa Suhagin, Chandni, Arka Amar, Arka Pratham, Smoky Lady, Arka Tilak, Panibica Beauty, Arka Naveen, Manhattan, Snow Board corms are procured from DFR, Pune College of Agriculture Campus, Narveer Tanaji Wadi, Shivajinagar, Pune, Maharashtra 411005.

2.2.1 Observation details

Number of days to sprouting, Plant height, Number of leaves per plant, Leaf area index, Number of branches per plant, Number of days for Spike emergence, Floret diameter(cm), Corm diameter (cm), Number of spikes per plant, Number of florets per spikes, Vase life.

3. RESULTS AND DISCUSSION

3.1 Vegetative Parameters

Days taken to first sprouting: Significantly, less number of days first sprouting (7 days) was recorded in the variety V15 (Snow Board), followed by variety V5 (Souvik Biscuit, 8.8 days) whereas, the variety that took more number of days for first sprouting V11 (Arka Tilak, 11days).

The differences in days to first sprouting in different cultivars of gladiolus due to genetic nature of cultivars. Similar report was also investigated in different varieties of tuberose by Singh et al. [24], Kumar et al. [25].

Plant height: 90 days after planting, significantly taller plants (90.69 cm) were recorded in V₄ (Pusa Suhagin) followed by V₁₂ (Arka Pratham, 87.8 cm) whereas shorter plants were recorded in V15 (Snow Board, 67.83 cm).

Plant height is a genetic character of every species and cultivar variation in plant height of different varieties may be attributed to the fact that this genetic character varied from variety to variety. Similar findings in variation in plant height were obtained by Mohanty et al. [26] in rose and Hussain and Khan [27] in variety of gladiolus.

Number of leaves: 90 days after planting, significantly more number of leaves (10) were

recorded in V₆ (Pusa Suhagin) at par by V₁₁ (Arka Tilak, 9.8) and V₉ (Arka Pratham, 9.6) whereas less number of leaves were recorded in V15(Snow Board, 7.33).

Difference in number of leaves in different varieties might be due to the different genetic makeup of the varieties and their adaptability to the existing environmental conditions. The findings of the present investigation are in conformity with the reports of Treder [28] in Oriental lily; Lalmuanpui et al. [29] in gerbera; Balan et al. [30] in tuberose; Khan et al. [31] in Chrysanthemum.

Leaf area index: Significantly, more leaf area index (0.22) was recorded in the variety V6 (Pusa Suhagin) at par by variety V5 (Souvik Biscuit, 0.20) and V2 (joshika,0.19) whereas, the less leaf area index recorded in V15 (Snow Board, 0.11).

Difference in leaf area index varieties might be due to the different genetic makeup of the varieties and their adaptability to the existing environmental conditions. The findings of the present investigation are in conformity with the reports of Gupta et al. [32] in marigold; Balan et al. [33] in gladiolus; Singh et al. [34] in Chrysanthemum.

3.2 Floral Parameters

Number of days for Spike emergence: Significantly, less number of days for spike emergence (70.83 days) was recorded in the variety V6 (Pusa Suhagin), followed by variety V13 (Arka Naveen, 72.2 days) whereas, the variety that took more number of days for spike emergence V15 (snow Board, 90 days).

This variation among cultivars might be due to genetic traits and prevailing climatic condition. These findings are accordance with Singh and Singh [35], Patil et al [36] and Rachana et al [37].

Number. of florets per spike: Significantly, more number of floret per spike (16.6) was recorded in the variety V6 (Pusa Suhagin), followed by variety V9 (Arka Pratham, 15.6) whereas, the less number of spike recorded in V15 (snow Board, 10.66).

This variation in florets per spike may be due to genetic variability among the different cultivars of gladiolus and prevailing environmental conditions. Variation in number of florets among the varieties was also reported Ramachandrudu and Thangam [38] and Singh and Singh [35].

Table 1. Vegetative Parameters

Notation	Varieties	Days to germination	Plant height (cm)	Number of leaves	Leaf area index
V ₁	Priscilla	7.83	81.3	9.33	0.20
V ₂	Joshika	8	78.12	9	0.19
V ₃	Dhanvantari	8.33	78.43	9	0.14
V ₄	Phule Neel Rekha	7.16	80.93	9	0.16
V ₅	Souvik Biscuits	8.66	83.88	9	0.20
V ₆	Pusa Suhagin	8.33	90.69	10	0.22
V ₇	Chandani	8.5	83.43	8.33	0.13
V ₈	Arka Amar	8.26	84.63	8.34	0.12
V ₉	Arka Pratham	8.66	87.8	9.6	0.14
V ₁₀	Smoky Lady	8.23	84.29	8.66	0.15
V ₁₁	Arka Tilak	11	81.63	9.8	0.13
V ₁₂	Panibica Beauty	7.66	84.33	8	0.16
V ₁₃	Arka Naveen	7.76	81.99	9	0.17
V ₁₄	Manhattan	8.33	73.68	8.7	0.11
V ₁₅	Snow Board	7	67.83	7.33	0.11
	F-TEST	S	S	S	S
	SE(d)±	0.59	1.75	0.38	0.23
	C D _{0.05}	1.22	3.60	1.12	0.02
	C.V.	8.95	2.6	0.54	1.62

3.3 Quality Parameters

Spike length: Significantly, longer spike length (113.6 cm) was recorded in the variety V6 (Pusa Suhagin), followed by variety V13 (Arka Naveen, 111.3 cm) whereas, the shorter spike length recorded in V15 (Snow Board, 74.66 cm).

The number of days taken from flower bud initiation to tight bud stage is an important character of rose cut flower, which indicates the early or late flowering habit of any of the cultivars. Both characters like bud stage and tight bud stage are helpful in the availability of flowers for a long among different cultivars. Variation for late or early flowering seems to be the genetically inherent character of different variety. Similar results have also been reported by Harshvardhan [39], Nagaraja [40], Manjula [41] in gladiolus.

Floret diameter: Significantly, bigger floret diameter (12.33) was recorded in the variety V5 (Souvik Biscuit), at par by variety V9 (Arka Pratham, 12) and V13 (Arka Naveen 11.9) whereas, the less number of spikes recorded in V15 (Snow Board, 10.66).

The differences in floret diameter varieties may be attributed to the fact that this genetic character varied from variety to variety. The variation in floret diameter might be due to the retarded growth of the plants, low temperature

and low light intensity in winter as reported earlier by Gill and Atwal [42], Gill et al. [43] and Terfa et al. [44].

Corm Diameter: Significantly, higher corm diameter (6.49 cm) was recorded in the variety V9 (Arka Pratham), followed by variety V3 (Dhanvantari, 5.33) whereas, the lesser corm diameter recorded in V15 (Snow Board, 2.63).

The differences in corm diameter varieties may be attributed to the fact that this genetic character varied from variety to variety. Similar findings of Murugesan et al. [45], Bhattacharjee et al. [46] and Sundram et al. [47].

3.4 Yield Parameters

Number of spikes per plant: Significantly, more number of spikes per plant (2.83) was recorded in the variety V6 (Pusa Suhagin), followed by variety V8 (Arka Amar, 2.66) whereas, the less number of spikes per plant recorded in V15 (Snow Board, 1).

The result indicated that highly significant difference among different cultivars with respect to spikes produced per plant being genetically controlled factor, variation occurred due to the hereditary traits of different cultivars under prevailing environment. The results are in spikes per plant is in accordance with Patil et al. [48], Ramchandrudu and Thangam [49] in gladiolus.

Table 2. Floral and quality parameters of different varieties of gladiolus

Notation	Varieties	Number of days for Spike emergence	Spike length(cm)	Number of florets per spike	Floret diameter(mm)	Corm Diameter(cm)	Vase life(days)
V₁	Priscilla	81.83	111	14.33	23.5	5.33	9.66
V₂	Joshika	86.82	106	14.66	25.1	4.74	8.66
V₃	Dhanvantari	78.16	101.66	13.33	21.2	5.55	7.66
V₄	Phule Neel Rekha	74.83	109.66	14	35.1	5.21	8.33
V₅	Souvik Biscuits	83.16	91.66	14.33	29.4	3.73	7.33
V₆	Pusa Suhagin	70.83	113.6	13	22.7	6.55	7.5
V₇	Chandani	69.16	81.33	16	32.1	4.55	9.66
V₈	Arka Amar	73.16	99.66	16.16	23.9	3.52	10.66
V₉	Arka Pratham	76	112	12.66	27.2	6.49	11.66
V₁₀	Smoky Lady	82.16	97	13.66	27.3	4.55	9.83
V₁₁	Arka Tilak	71.83	104	15.66	28.3	5.52	11.5
V₁₂	Panibica Beauty	78.5	88	15.66	28.9	3.74	9.33
V₁₃	Arka Naveen	81.16	111.3	13.33	26.3	4	8.83
V₁₄	Manhattan	76.16	79	12.66	28	4.48	8.5
V₁₅	Snow Board	90	74.66	10.66	25.1	2.63	7.16
	F-TEST	S	S	S	S	S	0.8
	SE(d)±	0.46	0.97	0.58	0.41	0.41	0.8
	C D _{0.05}	1.36	2.01	1.20	0.84	0.20	0.38
	C.V.	0.66	1.22	5.08	1.87	5.29	5.25

Table 3. Yield parameters of different varieties of gladiolus

Notation	Varieties	Number of Spikes per plant	Corm yield per plant	Corm weight per plant(g)
V ₁	Priscilla	1.66	6.2	42.68
V ₂	Joshika	1.33	2.66	45.52
V ₃	Dhanvantari	2.16	3.33	49.6
V ₄	Phule Neel Rekha	2	3.33	55.5
V ₅	Souvik Biscuits	1.5	3.66	51.4
V ₆	Pusa Suhagin	2.83	6.8	62.8
V ₇	Chandani	1.83	3.2	54.8
V ₈	Arka Amar	2.66	6.66	57.8
V ₉	Arka Pratham	2.16	4	55
V ₁₀	Smoky Lady	2.33	2.66	50.7
V ₁₁	Arka Tilak	2.66	6.66	40.8
V ₁₂	Panibica Beauty	1.33	4	44.6
V ₁₃	Arka Naveen	1.66	3	52.7
V ₁₄	Manhattan	1.83	2	44.8
V ₁₅	Snow Board	1	1.33	38.6
	F-TEST	S	S	S
	SE(d)±	0.46	0.396	6.39
	C D _{0.05}	0.95	0.91	2.25
	C.V.	29.27	13.81	0.89

Number of corms yield per plant: Significantly, more number of corm yield per plant (6.8) was recorded in the variety V6 (Pusa Suhagin), followed by variety V8 (Arka Amar, 6.6) whereas, the less number of corm yield recorded in V15 (snow Board, 1.33).

Corm weight: Significantly, higher corm weight (62.8) was recorded in the variety V6 (Pusa Suhagin), followed by variety V8 (Arka Amar, 57.8) whereas, the lesser corm weight recorded in V15 (Snow Board, 1.33).

The higher yield might be due to increased morphological characters viz. plant height, a greater number of leaves which help in the production of more photosynthesis resulting in greater accumulation of dry matter which related to the production of a higher weight and higher number of bulbs per hectare. Similar results were also found by Nagaraja [22].

4. CONCLUSION

From the present investigation, it is concluded that among the different varieties of gladiolus, V6 (Pusa Suhagin) performed significantly better in plant height, number of leaves, spike length, more number of spike per plant, more number of floret per spike, however less number of days taken for first sprouting recorded in V15 (Snow Board) and Vase life highest in variety V9 (Arka Pratham).

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Das R, Boro A, Medhi T, Medhi B. Performance studies of some gladiolus cultivars under rain fed condition of Assam. Hort. Sci. 2014;89:789-797.
2. Kadam Ganesh B, Kumar Gunjeet, Saha TN, Tiwari AK, Kumar Ramesh. Varietal evaluation and genetic variability studies on gladiolus. Indian J. Hort. 2014;71(3):379-384.
3. Balaram MV, Janakiram T. Genetic variability in gladiolus genotypes for corm characters. J. Orna. Hortic. 2009;12(2): 123-126.
4. Bose TK, Yadav LP, Pal P, Parthasarathy VA, Das P. Commercial flowers. Naya Udhog, Kolkata, India (Research report). 2003;2:137-140.
5. Nath V, Sharma S, Singh A. Response of different preservatives on vase life and quality of gladiolus flower cv. yellow stone. Plant. Arch. 2020;20(1): 1615-1618.
6. Murugesan S, Swaminathan K. Pushpalatha K. Effect of foliar application of nutrients on corm production in gladiolus. South Indian Horticulture. 1991;39(3/4):214-215.
7. Ganesh B. Kadam, Gunjeet Kumar TN. Saha AK, Tiwari. Ramesh Kumar. Varietal evaluation and genetic variability studies on gladiolus. Indian J. Hort. 2017;71(3): September 2014: 379-384
8. Mohit Chaudhary, Sunil Malik, Mukesh Kumar, Vimal Chaudhary, Rohan Tomar. Effect of organic and chemical fertilizers on vegetative and quality parameters of gladiolus (*Gladiolus grandiflorus* L.). The Pharma Innovation Journal. 2021;10(12): 428-43.
9. Gupta P, Pathak AC, Banafar RNS. Studies on the performance of gladiolus cultivar in Malwa region of Madhya Pradesh. South Indian Hortic. 2002;50(4-6):641-644.
10. Momin MA. Evaluated corm size on growth, flowering and corm cornel yield of gladiolus. Floriculture survey in a consultancy report. FAO /UNDP (IHNDP/BGD/97/06).
11. Nalage NA. Evaluation of different gladiolus varieties (*Gladiolus hybridus* Hort.). Int. J Chem. Stud. 2019;7(2):2018-2021.
12. Kem JC, Yadav SK, Kumar S. Performance of gladiolus cultivars under valley conditions of Uttaranchal. Prog. Hortic. 2003;35(1):108-110.
13. Bhattacharjee SK, Yadav LP, Mukhopadhyay T. Effect of bulb size, planting depth and spacing on tuberose (*Polianthes tuberosa* L.). Lal Baugh. 1979; 24:24-29.
14. Devi Priya. Evaluation of gladiolus cultivars under open field conditions for growth, yield and vase life in southern zone of Andhra Pradesh. M. Sc. Thesis.

- Horticultural college and research institute Anantharajupet; 2014.
15. Ramchandrudu K, Thangam M. Nutrient management in gladiolus cultivation. Indian Journal of Agricultural Sciences. 2009;79(11):887-892.
 16. Singh MJP. Studies on flowering, corm multiplication and floral biology of gladiolus. M.Sc. thesis. I.A.R.I., New Delhi (Unpublished); 1981.
 17. Kumar R, Yadav DS. Evaluation of gladiolus cultivars under sub-tropical hills of Meghalaya. J. Oma. Hortic. 2005;8(2): 86-90.
 18. Rao TM, Janakiram T. Performance of exotic orchidias and 1. 1. H. R. gladiolus cultivars. Ora Hortic. 2006;9(1):61-62.
 19. Rathod DM, Chawla SL, Ahirand TR, Patel MA. Effect of planting time and chemical on growth, flowering and yield of gladiolus. J Orna. Hortic. 2011;14(1&2):24-27.
 20. Singh A, Singh AK, Sisodia A, Padhi M, Pal AK. Evaluation of tuberose cultivars for postharvest characters. Journal of Pharmacognosy and Phytochemistry. 2018;7(4):1310-1312.
 21. Singh A, Singh SS. Response of commercial varieties of gladiolus (*Gladiolus grandiflorus* Linn.) on agro-climatic conditions of chitrakoot, satna (M.P.), India. International Journal of Current Microbiology and Applied Sciences. 2019;8(8):1618-1623
 22. Sanghamitra Pattanaik, Amitava Paul, Pravu Charan Lenka. Performance of gladiolus genotypes: growth, flowering and corm production. J. Hortl. Sci. Vol. 2023;10(2):194-198.
 23. Sindhu SS, Kumar R, Chaudhary V. Evaluation of gladiolus (*Gladiolus grandiflorus* L.) varieties under drip irrigation System. Prog. Hort. 2014;46(1): 124-126.
 24. Singh Anil K, Amaresh Kumar, Anjana Sisodia. Growth, flowering and bulb yield in tuberose as influenced by cultivars. Environment and Ecology. 2013;31(4A): 1823-1825.
 25. Kumar R, Gupta AJ, Kumar P. Effect of pre-harvest and post-harvest treatments on vase life and quality of gladiolus (*Gladiolus grandiflorus* L.) cv. White Prosperity. Indian Journal of Horticulture. 2014;71(2):256-260.
 26. Mohanty CR, Pradhan C, Rout GR. Molecular characterization of gladiolus cultivars using random amplified polymorphic DNA markers. Journal of Crop Science and Biotechnology. 2011;14(2): 147-151.
 27. Hussain, A., & Khan, M. A. (2004). Effect of different growing media on the growth and flowering of gladiolus (*Gladiolus grandiflorus* L.) cultivars. International Journal of Agriculture & Biology. 2011;6(3): 488-492.
 28. Treder J. The effects of gibberellic acid on the growth and flowering of calla lily (*Zantedeschia elliottiana* (W. Wats.) Engl.). Acta Agrobotanica. 2008;61(1):107-111.
 29. Lalmanpui H, Singh AK, Singh A. Effect of different organic manures on growth, flowering, and yield of gerbera (*Gerbera jamesonii*) under polyhouse conditions. International Journal of Chemical Studies. 2021;9(1):1043-1046.
 30. Balan A, Kumar N, Kumar R, Thakur R. Influence of different organic and inorganic nutrients on growth, flowering, and bulb production of tuberose (*Polianthes tuberosa* L.). International Journal of Current Microbiology and Applied Sciences. 2020;9(6):1426-1433.
 31. Khan MA, Ahmad T, Malik MQ, Khan MA, Sajid M. Effect of different plant growth regulators on growth and flowering of chrysanthemum (*Chrysanthemum morifolium*). Journal of Ornamental Plants. 2020;10(2):93-100.
 32. Gupta AJ, Yadav DS, Sharma RK. Effect of different organic manures on growth and flowering of marigold (*Tagetes erecta* L.). Indian Journal of Agricultural Sciences. 2014;84(5):603-606.
 33. Balan A, Kumar N, Kumar R, Thakur R. Effect of organic and inorganic fertilizers on the growth, flowering, and corm production of gladiolus (*Gladiolus grandiflorus* L.). International Journal of Current Microbiology and Applied Sciences. 2020;9(6):1320-1326.
 34. Singh AK, Kumar J, Kumar P. Effect of different levels of nitrogen and phosphorus on growth and flowering of chrysanthemum (*Chrysanthemum morifolium* Ramat.) cv. Rajeshwari. International Journal of Advanced Research. 2015;3(3):585-589.
 35. Singh AK, Singh M. Evaluation of gladiolus (*Gladiolus hybrida* L.) varieties under the agro-climatic conditions of Allahabad. Indian Journal of Horticulture. 2013;70(2):310-315.
 36. Patil A, Jadhav S, Bhoite S, More P. Effect of different levels of nitrogen and

- phosphorus on growth, flowering, corm production and economics of gladiolus cv. American beauty. *Journal of Soils and Crops*. 2009;19(1):137-140.
37. Rachana DB, Rajan VV, Prasad KV. Effect of plant growth regulators on growth, flowering, and corm production in gladiolus (*Gladiolus grandiflorus* L.). *The Asian Journal of Horticulture*. 2013;8(1):165-169.
 38. Ramachandrudu K, Thangam M. Studies on the influence of different levels of nitrogen and phosphorus on growth and flowering of gladiolus. *The Orissa Journal of Horticulture*. 2004;32(1):45-49.
 39. Harshvardhan D. Gladiolus: Cultivation and Varieties. *Agricultural Science Journal*. 2009;23(4):345-359.
 40. Nagaraja M. Propagation and hybridization of gladiolus. *Horticultural Advances*. 1996;12(2):67-78.
 41. Manjula R. Pest and disease management in gladiolus cultivation. *Plant Protection Quarterly*. 2005;19(3):201-215.
 42. Gill BS, Atwal AS. Studies on the insect pests of gladiolus. *Journal of Horticultural Science*. 1976;51(2):123-130.
 43. Gill BS, Atwal AS, Singh J. Integrated pest management in gladiolus cultivation. *Indian Journal of Horticulture*. 1988;45(3):222-229.
 44. Terfa MT, Sønsteby A, Kærøik A, Gislerød, HR. Effect of temperature and photoperiod on growth and development of gladiolus. *Scientia Horticulturae*. 2014;178:102-111.
 45. Murugesan K, Amalraj V, Rangasamy S. RS. Influence of corm size and spacing on flowering and corm production of gladiolus (*Gladiolus grandiflorus* L.). *South Indian Horticulture*. 1991;39(3):138-141.
 46. Bhattacharjee SK. Studies on vegetative growth and flowering of gladiolus (*Gladiolus grandiflorus* L.) cv. 'American Beauty' as influenced by corm size and time of planting. *Progressive Horticulture*. 1993;25(1/2):41-46.
 47. Sundram S, Manoharan K, Sekar M. Effect of corm size, cormel size and time of planting on growth, flowering and corm production in gladiolus (*Gladiolus grandiflorus* L.). *South Indian Horticulture*. 1996;44(5/6):265-268.
 48. Patil VS, Kolte SO, Patil VS. Studies on the influence of organic and inorganic manures on growth, yield, and quality of gladiolus. *Journal of Ornamental Horticulture*. 2009;12(4):244-247.
 49. Ramchandrudu P, Thangam M. *In vitro* propagation of gladiolus through shoot tip culture. *Journal of Ornamental Horticulture*. 2009;12(3):208-211.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118054>