



G and P Genotypes of Group A Rotavirus in Diarrheic Calves in Costa Rica

José Bonilla-Espinoza^{1,2}, Lurys Bourdett-Stanziola^{1,3,4*} and Carlos Jiménez¹

¹*Programa de Investigación en Enfermedades Tropicales (PIET), Escuela de Medicina Veterinaria, Heredia, Costa Rica.*

²*Universidad Nacional Autónoma de Nicaragua, (UNAN), León, Nicaragua.*

³*Instituto de Investigaciones Científicas, INDICASAT, Panamá.*

⁴*Facultad de Ciencias Agropecuarias, Universidad de Panamá, Panamá.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Hazem Mohammed EbraheemShaheen, Damanhour University, Egypt.

Reviewers:

(1) Nawar Alsalih, Al-muthanna University, Iraq.

(2) Alexander Zakhartchouk, University of Saskatchewan, Canada.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here: <https://www.sdiarticle5.com/review-history/77547>

Short Communication

**Received 02 October 2021
Accepted 06 December 2021
Published 09 December 2021**

ABSTRACT

Rotavirus infections are the most common causes of infectious diarrhea in young children and animals worldwide. We performed a genotyping of rotavirus strains by RT-PCR for G (VP7) and P (VP4), in diarrhea samples from calves, in several locations in Costa Rica. Combinations of genotypes G (VP7) and P (VP4) were identified in cattle: 4 (26%) corresponding to G8 P [11], 4 (24%) to G6P [11], 3 (19%) to G8P [7], 2 (13%) to G10P [1], 3 (19%) to G10P [11], and 1 (6%) to G8P [1] and mixed infections. This research represents the first report of bovine rotavirus strains (calves) in Costa Rica and in the Central American region.

Keywords: Rotavirus; diarrhea; calves; VP7 and VP4.

1. INTRODUCTION

Rotavirus is considered the main etiological agent in diarrhea in young animals and children around the world [1].

Group A Rotaviruses are classified using a binary system, according to the genetic and antigenic variation of two surface proteins (VP7 and VP4). So far, 36 types G (VP7) and at least 51 types P (VP4) of rotavirus group A circulating in humans and animals [2] have been reported. Particularly in cattle, 14 different G (VP7) genotypes (G1 -G6, G8, G10-G12, G15, G18, G21, and G24 and 11 P (VP4) genotypes (P [1], P [3], P [5] -P [7], P [11], P [14], P [17], P [21], P [29], and P [33]) [3,4] have been identified. They affect animal populations causing great economic losses for livestock producers, due to vaccination expenses, treatment, weight loss of affected animals and mortality.

Risk factors that contribute to the appearance of diarrhea caused by rotavirus are failures in the transfer of passive immunity especially observed in heifers, presence of seasonal diarrhea, and lack of hygiene in artificial rearing systems [5].

Knowledge of the epidemiology of rotavirus is essential for the control of this disease; especially since it has been observed that it presents differences between developed and developing countries, as well as differences in climate [6].

In Costa Rica, in the Central American region and in the Dominican Republic we have carried out characterizations of rotavirus strains in children, where we reported common and unusual genotypes [7].

This research represents the first genotyping study of rotavirus G (VP7) and P (VP4) strains detected in cattle in Costa Rica and in the Central American region.

2. STUDY DESCRIPTION

Samples of feces from calves under 1 month of age were taken from 17 dairy farms (bovine-calves) located in Río Cuarto de Grecia, Pital, Pos, Zarcero, Sucre and Pacayas in 2008 in Costa Rica. All calf stool samples were analyzed in the virology laboratory, at the School of Veterinary Medicine at the National University of Costa Rica (UNA), to determine the presence of rotavirus using the immunochromatography technique (CorisBioConcept, Belgium) and the RT-PCR technique (VP7) and (VP4) described by Genstch JR et al [8] and Gouvea V et al [9].

From 200 samples (calves feces), a total of 38 samples were ELISA positive for rotavirus and 25 samples were sent for characterization of the G (VP7) and P (VP4) genotype.

By using the RT-PCR technique, the combinations of genotypes G (VP7) and P (VP4) were identified in cattle: 4 (26%) corresponding to G8 P [11], 4 (24%) to G6 P [11], 3 (19%) to G8 P [7], 2 (13%) to G10 P [1], 3 (19%) to G10P [11], and 1 (6%) to G8P [1]. In 3 samples, it was not possible to identify the G (VP7) genotype with the primers used. Mixed infections were found in 6 samples (Table 1).

The distribution of genotypes G (VP7) and P (VP4) of Group A bovine rotavirus in the dairy farms of the different locations in Costa Rica, were different. Namely, G10 was the most prevalent genotype in the Poas, Zarcero and Sucre area, while the G6 genotype prevailed in the Río Cuarto de Grecia, Poas and Pital de San Carlos areas. Finally, G8 was identified mainly in the Pacayas, Poas, Cedral de San Carlos and Zarcero areas. In the P (VP4) types, P [11] was identified as the main genotype in Río Cuarto de Grecia, Pital de San Carlos, Poas, Cedral de San Carlos and Pacayas, P [7] was found only in Río Cuarto de Grecia, whereas P [1] was identified in Zarcero and Sucre. Mixed infections were identified in Pacayas and Poas (Fig. 1) (Table 2).

Table 1. Distribution of G (VP7) P (VP4) of rotavirus strains isolated from bovine in Costa Rica

Genotype	P [11]	P[11]+P[7]	P[7]	P[1]	P[11]+P[1]	Total
G8	2(13)	2(13)	3(19)	1 (6)	0	8(51)
G6	4(24)	0	0	0	0	4(24)
G10	2(13)	0	0	1(6)	1(6)	4(25)
Total	8(50)	2(13)	3(19)	2 (12)	1(6)	16 (100)

No. (%) positive (n=16)




Fig. 1. Distribution of Genotypes G (VP7) and P (VP4) in Sampled Localities of Costa Rica

Table 2. Distribution of genotypes G (VP7) and P (VP4) of Group A Bovine Rotavirus in 17 dairy farms sampled in the year 2008

Farm	Location	Samples Collected	Positive Samples	G (VP7)	P (VP4)
1	Río Cuarto de Grecia	22	6	G6	P[11]
2	Poas	21	7	G6 , G10	P[11]
3	Pital San Carlos	11	2	G6	P[11]
4	Poas	13	0		
5	Poas	8	1	G8	
6	Poas	7	0		
7	Poas	9	0		
8	Poas	8	0		
9	Poas	7	2	G10	P[11], P[1]
10	Poas	10	1		P[11], P[1]
11	Poas	8	0		
12	Pacayas, Cartago	18	4	G8	P[11], P[7]
13	Cedral, San Carlos	5	1	G8	P[11]
14	Zarcero	10	1	G8	P[1]
15	Sucre	16	1	G10	P[1]
16	Zarcero	12	1	G10	
17	Sucre	15	1	G10	
Total		200	28		

There are reports in the literature of infections in cattle caused by various genotypes of Group A Rotavirus such as G1, G2, G3, G6, G8, G10 and G11 [4,10-12]. In the present study, the (VP7)

G6, G8 and G10 genotypes were identified, with the G10 genotype prevailing, which was identified in 10 samples, followed by G6 and G8 identified in 8 and 7 fecal samples, respectively.

It is important to clarify that the farms sampled in Costa Rica all belonged to the province of Alajuela except for Pacayas, which belongs to the province of Cartago, both provinces are part of the Central Valley Mountain Range of the country. Our findings coincide with reports in Brazil, Argentina, and other countries in the world where high frequencies of these genotypes were reported and considered common of epidemiological importance in cattle [10-12]. Regarding the genotypes P (VP4), we have the P [1], P [5] and P [11] as frequent as they reported worldwide [10-12]. In this work, the genotype P [11] was the most frequent, followed by P [1] that was identified in 2 positive samples and P [7]. In addition, mixed infections corresponding to P [11 + P1] and P [11 + P7] were identified. For example, Da Silva Medeiros et al [10], found P [11] as the most frequent genotype in calf diarrhea in Brazil. The finding of mixed infections is very important from the epidemiological point of view, since animals affected by one type of strain can be infected with another different strain or be affected by two different strains at the same time, which makes control of the illness even more difficult.

Among the findings from different locations in Costa Rica, a variation of P groups (VP4) was detected, but despite the variety of P groups found in different areas of the country, they are similar to those described in other parts of the world [11-13]. Additionally, as previously mentioned, the locations sampled belong to the Central Valley of Costa Rica, for which it would be important to carry out another study in which more samples are collected from other regions of the country.

Within the different combinations of genotypes G (VP7) and P (VP4) of Group A Rotavirus in bovines, G6 P [11] was identified as the most frequent detected in 4 of the 16 positive samples, G8 P [7] was the second most frequent combination identified in 3 of the 16 positive samples, followed by G8 P [11] and G10 P [11] found in 2 samples; G8 P [1] and G10 P [1] were identified in one sample. Our findings differ partially from the combinations of G6 P genotypes [5] reported as the most frequent in South America [10,11]. Studies carried out in children have also revealed the presence of the G8, G10 genotype in America and other continents [14,15,16]. Although we also identified mixed infections in this investigation; studies have shown that rotaviruses produce

“reassortants”. These genetic changes contribute to the evolution of rotaviruses, resulting in a diversity of genotypes and mixed infections, which is characteristic of rotavirus infections [1,3,13,14]. A literature review reveals the emergence of new strains of rotavirus in the American continent [14]. Studies made on the phylogenetics of rotavirus have demonstrated differences in the lineage and sub lineage of genotypes, highlighting the great genetic variability of rotavirus [1]. A small percentage of samples analyzed in this study could not be typed with the use of specific primers, which may be due to the accumulation of point mutations and diversity of reported lineages. We consider that additional experiments and sequencing analysis of these genotypes should be carried out in future investigations, because studies carried out in Nicaragua, Guatemala, Honduras, and the Dominican Republic have reported the presence of rotavirus strains of bovine origin, pigs, and bats in rotavirus samples, excreted from children with diarrhea, showing possible zoonosis [17-20]. Furthermore, studies carried out in Uruguay have detected human rotavirus genes in calf diarrhea samples [13]. There is a report on the detection analysis of rotavirus strains in bats in Costa Rica. This investigation also included taking samples of bat feces from other countries in the world, revealing strains of rotavirus never reported before, even some of them have similarities with rotavirus strains found in other mammals [21]. Currently, commercial vaccines are available for the prevention of neonatal diarrhea in calves and children that have shown varying degrees of effectiveness. These vaccines are manufactured with the genotypes most frequently found and reported in different parts of the world. In this context, the immunogenicity and efficacy of rotavirus vaccines may be challenged by the evolution of the rotavirus viral genome.

3. CONCLUSION

Therefore, and adding that research reveals that rotaviruses cause deaths in children, cattle and pigs, we consider urgent the need to maintain an epidemiological surveillance of rotavirus strains in children, domestic animals, wild animals, cattle and pigs in the country, as well as the reinforcement in the maintenance and consolidation in the unification of detection and characterization protocols of rotavirus in laboratories in Central America.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

ACKNOWLEDGEMENTS

This study was supported by the Network for Research and Training of Tropical Diseases in Central America (NeTropica), (Grant: 2005-07). We thank our Tutor Dr. Carlos Jiménez for all his guidance and support in this research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Luchs A, Timenetsky Mdo C. Group a Rotavirus gastroenteritis: Post-vaccine era, genotypes and zoonotic transmission. *Einstein (Sao Paulo)*. 2016;14(2):278-87.
2. RCWG, Rotavirus Classification Working Group, Newly Assigned Genotypes - Update May 29th 2018, (Accessed 2 July 2018), Available:<https://rega.kuleuven.be/cev/viraImetagenomics/virus-classification/rcwg>.
3. Martella V, Bányai K, Matthijnssens J, Buonavoglia C, Ciarlet M. Zoonotic aspects of rotaviruses. *VetMicrobiol*. 2010; 140(3-4):246-55.
4. Matthijnssens J, Ciarlet M, McDonald SM, Attoui H, Bányai K, Brister JR, et al. Uniformity of rotavirus strain nomenclature proposed by the Rotavirus Classification Working Group (RCWG). *Arch Virol*. 2011; 156(8):1397-413.
5. Teodoroff T, Tsunemitsu H, Okamoto K, Katsuda K, Kohmoto M, Kawashima K, et al. Predominance of porcine rotavirus G9 in Japanese piglets with diarrhea: close relationship of their VP7 genes with those of recent human G9 strains. *Journal of Clinical Microbiology*. 2005;43(3):1377-1384.
6. De Oliveira LH, Danovaro-Holliday MC, Andrus JK, de Fillipis AM, Gentsch J, Matus CR, Widdowson MA; Rotavirus Surveillance Network. Sentinel hospital surveillance for rotavirus in Latin American and Caribbean countries. *J Infect Dis*. 2009;200 Suppl 1:S131-9.
7. Bourdett-Stanziola L, Ortega-Barria E, Espinoza F, Bucardo F, Jimenez C, Ferrera A. Rotavirus genotypes in Costa Rica, Nicaragua, Honduras and the Dominican Republic. *Intervirology*. 2011; 54(1):49-52.
8. Gentsch JR, Glass RI, Woods P, Gouvea V, Gorziglia M, Flores J, et al. Identification of group A rotavirus gene 4 types by polymerase chain reaction. *J ClinMicrobiol*. 1992;30(6):1365-73.
9. Gouvea V, Santos N, TimenetskyMdo C. Identification of bovine and porcine rotavirus G types by PCR. *J ClinMicrobiol*. 1994;32(5):1338-40.
10. da Silva Medeiros TN, Lorenzetti E, Alfieri AF, Alfieri AA. G and P genotype profiles of rotavirus A field strains circulating in beef and dairy cattle herds in Brazil, 2006-2015. *Comp ImmunolMicrobiol Infect Dis*. 2019;64:90-98.
11. Badaracco A, Garaicoechea L, Rodríguez D, Uriarte EL, Odeón A, Bilbao G, et al. Bovine rotavirus strains circulating in beef and dairy herds in Argentina from 2004 to 2010. *Vet Microbiol*. 2012;158(3-4):394-9. A.
12. Cashman O, Lennon G, Sleator RD, Power E, Fanning S, O'Shea H. Changing profile of the bovine rotavirus G6 population in the south of Ireland from 2002 to 2009. *Vet Microbiol*. 2010;146(3-4):238-44.
13. Castells M, Caffarena RD, Casaux ML, Schild C, Miño S, Castells F, et al. Phylogenetic Analyses of Rotavirus A from Cattle in Uruguay Reveal the Circulation of Common and Uncommon Genotypes and Suggest Interspecies Transmission. *Pathogens*. 2020;9(7):570.
14. Bourdett-Stanziola L, Centeno E, Cuevas-Abrego M, Durant-Archibold A. A, Ortega-Barria E, Bucardo F. The Emergence of New Rotavirus Strains in America. *Review. South Asian Journal of Research in Microbiology*. 2021;11(1):46-62. DOI.org/10.9734/sajrm/2021/v11i130244.
15. Esona MD, Page NA, Akran VA, Armah GE, Steele AD. Characterization of 2 human genotype G10 rotavirus strains, 3008CM and 1784/CI/1999, isolated in

- Cameroon and Cote d'Ivoire during the 1999-2000 rotavirus season. *J Infect Dis.* 2010;202Suppl:S212-9.
16. Zuccotti G, Meneghin F, Dilillo D, Romano L, Bottone R, Mantegazza C, et al. Epidemiological and clinical features of rotavirus among children younger than 5 years of age hospitalized with acute gastroenteritis in Northern Italy. *BMC Infect Dis.* 2010;10:218.
 17. Bányai K, Esona MD, Mijatovic S, Kerin TK, Pedreira C, Mercado J, et al. Zoonotic bovine rotavirus strain in a diarrheic child, Nicaragua. *J Clin Virol.* 2009;46(4): 391-3.
 18. Quaye O, Roy S, Rungsisuriyachai K, Esona MD, Xu Z, Tam KI, et al. Characterization of a rare, reassortant human G10 P [14] rotavirus strain detected in Honduras. *Mem Inst Oswaldo Cruz.* 2018;113(1):9-16.
 19. Gautam R, Mijatovic-Rustempasic S, Roy S, Esona MD, Lopez B, Mencos Y, et al. Full genomic characterization and phylogenetic analysis of a zoonotic human G8P [14] rotavirus strain detected in a sample from Guatemala. *Infect Genet Evol.* 2015;33:206-11.
 20. Bourdett-Stanziola L, Centeno E, Nordgren J, Durant-Archibold A, Ortega-Barría E and Bucardo F. Potential Bat-like rotavirus in Hospitalized Children with Diarrhea from the Dominican Republic. *Am J Res Infect Dis.* 2021; 8(1):1-7. DOI.org/10.9734/ajrid/2021/v8i130225.
 21. Simsek C, Corman VM, Everling HU, Lukashev AN, Rasche A, Maganga GD, et al. At Least Seven Distinct Rotavirus Genotype Constellations in Bats with Evidence of Reassortment and Zoonotic Transmissions. *mBio.* 2021;12(1):e02755-20.

© 2021 Bonilla-Espinoza et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77547>*