



Parameter Estimation of Length Biased Weighted Frechet Distribution via Bayesian Approach

Arun Kumar Rao¹ and Himanshu Pandey^{1*}

¹Department of Mathematics and Statistics, DDU Gorakhpur University, Gorakhpur, India.

Authors' contributions

This work was carried out in collaboration between both authors. Author AKR designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Author HP managed the analyses of the study and literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJPAS/2021/v11i130259

Editor(s):

(1) Dr. Thomas L. Toulas, University of West Attica, Greece.

Reviewers:

(1) Juan Pablo Molina Aguilar, Universidad Michoacana de San Nicolás de Hidalgo (UMSNH), Mexico.

(2) Gannat Ramadan AL-Dayian, Al-Azhar University, Egypt.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/65210>

Received: 02 December 2020

Accepted: 09 February 2021

Published: 24 February 2021

Original Research Article

Abstract

In this paper, length-biased weighted Frechet distribution is considered for Bayesian analysis. The expressions for Bayes estimators of the parameter have been derived under squared error, precautionary, entropy, K-loss, and Al-Bayyati's loss functions by using quasi and gamma priors.

Keywords: Bayesian method; length-biased weighted frechet distribution; quasi and gamma priors; squared error; precautionary; entropy; k-loss and al-bayyati's loss functions.

1 Introduction

Reyad et al. [1] was introduced a new class of length-biased weighted Frechet distribution. They obtained the statistical properties of this distribution and estimated the parameters by using moment and likelihood approaches. The main objective of this paper is to obtain the Bayes estimators of the parameter under different loss functions by using quasi and gamma priors. The probability density function (pdf) of length-biased weighted Frechet distribution is given by

*Corresponding author: E-mail: himanshu_pandey62@yahoo.com;

$$f(x; \theta) = \frac{a}{\Gamma\left(1 - \frac{1}{a}\right)} \theta^{\left(1 - \frac{1}{a}\right)} x^{-a} e^{-\theta x^{-a}} \quad ; x > 0, a > 1. \quad (1)$$

The joint density function or likelihood function of (1) is given by

$$f(\underline{x}; \theta) = \left(a / \Gamma\left(1 - \frac{1}{a}\right) \right)^n \theta^{n\left(1 - \frac{1}{a}\right)} \left(\prod_{i=1}^n x_i^{-a} \right) e^{-\theta \sum_{i=1}^n x_i^{-a}}. \quad (2)$$

The log likelihood function is given by

$$\log f(\underline{x}; \theta) = n \log \left(a / \Gamma\left(1 - \frac{1}{a}\right) \right) + n \left(1 - \frac{1}{a}\right) \log \theta + \log \left(\prod_{i=1}^n x_i^{-a} \right) - \theta \sum_{i=1}^n x_i^{-a}. \quad (3)$$

Differentiating (3) with respect to θ and equating to zero, we get the maximum likelihood (ML) estimator of θ which is given as

$$\hat{\theta} = n \left(1 - \frac{1}{a}\right) \left(\sum_{i=1}^n x_i^{-a} \right)^{-1}. \quad (4)$$

2 Bayesian Method of Estimation

The Bayesian inference procedures have been developed generally under squared error loss function

$$L(\hat{\theta}, \theta) = (\hat{\theta} - \theta)^2. \quad (5)$$

The Bayes estimator under the above loss function, say, $\hat{\theta}_s$ is the posterior mean, i.e,

$$\hat{\theta}_s = E(\theta). \quad (6)$$

Zellner [2], Basu and Ebrahimi [3] have recognized that the inappropriateness of using symmetric loss function. Norstrom [4] introduced precautionary loss function is given as

$$L(\hat{\theta}, \theta) = \frac{(\hat{\theta} - \theta)^2}{\hat{\theta}}. \quad (7)$$

The Bayes estimator under this loss function is denoted by $\hat{\theta}_P$ and is obtained as

$$\hat{\theta}_P = [E(\theta^2)]^{1/2}. \quad (8)$$

Calabria and Pulcini [5] points out that a useful asymmetric loss function is the entropy loss

$$L(\delta) \propto [\delta^p - p \log_e(\delta) - 1]$$

where $\delta = \frac{\hat{\theta}}{\theta}$, and whose minimum occurs at $\hat{\theta} = \theta$. Also, the loss function $L(\delta)$ has been used in Dey et al. [6] and Dey and Liu [7], in the original form having $p = 1$. Thus $L(\delta)$ can written be as

$$L(\delta) = b[\delta - \log_e(\delta) - 1]; \quad b > 0. \tag{9}$$

The Bayes estimator under entropy loss function is denoted by $\hat{\theta}_E$ and is obtained by solving the following equation

$$\hat{\theta}_E = \left[E\left(\frac{1}{\theta}\right) \right]^{-1}. \tag{10}$$

Wasan [8] proposed the K-loss function which is given as

$$L(\hat{\theta}, \theta) = \frac{(\hat{\theta} - \theta)^2}{\hat{\theta}\theta}. \tag{11}$$

Under K-loss function the Bayes estimator of θ is denoted by $\hat{\theta}_K$ and is obtained as

$$\hat{\theta}_K = \left[\frac{E(\theta)}{E(1/\theta)} \right]^{\frac{1}{2}}. \tag{12}$$

Al-Bayyati [9] introduced a new loss function which is given as

$$L(\hat{\theta}, \theta) = \theta^c (\hat{\theta} - \theta)^2. \tag{13}$$

Under Al-Bayyati's loss function the Bayes estimator of θ is denoted by $\hat{\theta}_{Al}$ and is obtained as

$$\hat{\theta}_{Al} = \frac{E(\theta^{c+1})}{E(\theta^c)}. \tag{14}$$

Let us consider two prior distributions of θ to obtain the Bayes estimators.

- (i) Quasi-prior: For the situation where we have no prior information about the parameter θ , we may use the quasi density as given by

$$g_1(\theta) = \frac{1}{\theta^d}; \theta > 0, d \geq 0, \tag{15}$$

where $d = 0$ leads to a diffuse prior and $d = 1$, a non-informative prior.

- (ii) Gamma prior: Generally, the gamma density is used as prior distribution of the parameter θ given by

$$g_2(\theta) = \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{\alpha-1} e^{-\beta\theta}; \theta > 0. \tag{16}$$

3 Posterior Density under $g_1(\theta)$

The posterior density of θ under $g_1(\theta)$, on using (2), is given by

$$\begin{aligned} f(\theta/\underline{x}) &= \frac{\left[\left(\frac{a}{\Gamma\left(1-\frac{1}{a}\right)} \right)^n \theta^{n\left(1-\frac{1}{a}\right)} \left(\prod_{i=1}^n x_i^{-a} \right) e^{-\theta \sum_{i=1}^n x_i^{-a}} \theta^{-d} \right]}{\int_0^\infty \left[\left(\frac{a}{\Gamma\left(1-\frac{1}{a}\right)} \right)^n \theta^{n\left(1-\frac{1}{a}\right)} \left(\prod_{i=1}^n x_i^{-a} \right) e^{-\theta \sum_{i=1}^n x_i^{-a}} \theta^{-d} \right] d\theta} = \frac{\theta^{n\left(1-\frac{1}{a}\right)-d} e^{-\theta \sum_{i=1}^n x_i^{-a}}}{\int_0^\infty \theta^{n\left(1-\frac{1}{a}\right)-d} e^{-\theta \sum_{i=1}^n x_i^{-a}} d\theta} \\ &= \frac{\left(\sum_{i=1}^n x_i^{-a} \right)^{n\left(1-\frac{1}{a}\right)-d+1}}{\Gamma\left(n\left(1-\frac{1}{a}\right) - d + 1 \right)} \theta^{n\left(1-\frac{1}{a}\right)-d} e^{-\theta \sum_{i=1}^n x_i^{-a}} \end{aligned} \tag{17}$$

Theorem 1. On using (17), we have

$$E(\theta^c) = \frac{\Gamma\left(n\left(1-\frac{1}{a}\right) - d + c + 1 \right)}{\Gamma\left(n\left(1-\frac{1}{a}\right) - d + 1 \right)} \left(\sum_{i=1}^n x_i^{-a} \right)^{-c}. \tag{18}$$

Proof. By definition,

$$E(\theta^c) = \int \theta^c f(\theta/\underline{x}) d\theta$$

$$\begin{aligned}
 &= \frac{\left(\sum_{i=1}^n x_i^{-a}\right)^{n\left(1-\frac{1}{a}\right)-d+1}}{\Gamma\left(n\left(1-\frac{1}{a}\right)-d+1\right)} \int_0^\infty \theta^{n\left(1-\frac{1}{a}\right)-d+c} e^{-\theta \sum_{i=1}^n x_i^{-a}} d\theta \\
 &= \frac{\left(\sum_{i=1}^n x_i^{-a}\right)^{n\left(1-\frac{1}{a}\right)-d+1} \Gamma\left(n\left(1-\frac{1}{a}\right)-d+c+1\right)}{\Gamma\left(n\left(1-\frac{1}{a}\right)-d+1\right) \left(\sum_{i=1}^n x_i^{-a}\right)^{n\left(1-\frac{1}{a}\right)-d+c+1}} \\
 &= \frac{\Gamma\left(n\left(1-\frac{1}{a}\right)-d+c+1\right)}{\Gamma\left(n\left(1-\frac{1}{a}\right)-d+1\right)} \left(\sum_{i=1}^n x_i^{-a}\right)^{-c}.
 \end{aligned}$$

From equation (18), for $c = 1$, we have

$$E(\theta) = \left(n\left(1-\frac{1}{a}\right)-d+1\right) \left(\sum_{i=1}^n x_i^{-a}\right)^{-1}. \tag{19}$$

From equation (18), for $c = 2$, we have

$$E(\theta^2) = \left[\left(n\left(1-\frac{1}{a}\right)-d+2\right)\left(n\left(1-\frac{1}{a}\right)-d+1\right)\right] \left[\sum_{i=1}^n x_i^{-a}\right]^{-2}. \tag{20}$$

From equation (18), for $c = -1$, we have

$$E\left(\frac{1}{\theta}\right) = \frac{1}{\left(n\left(1-\frac{1}{a}\right)-d\right)} \sum_{i=1}^n x_i^{-a}. \tag{21}$$

From equation (18), for $c = c + 1$, we have

$$E(\theta^{c+1}) = \frac{\Gamma\left(n\left(1-\frac{1}{a}\right)-d+c+2\right)}{\Gamma\left(n\left(1-\frac{1}{a}\right)-d+1\right)} \left(\sum_{i=1}^n x_i^{-a}\right)^{-(c+1)}. \tag{22}$$

4 Bayes Estimators Under $g_1(\theta)$

From equation (6), on using (19), the Bayes estimator of θ under squared error loss function is given by

$$\hat{\theta}_S = \left(n \left(1 - \frac{1}{a} \right) - d + 1 \right) \left(\sum_{i=1}^n x_i^{-a} \right)^{-1}. \quad (23)$$

From equation (8), on using (20), the Bayes estimator of θ under precautionary loss function is obtained as

$$\hat{\theta}_P = \left[\left(n \left(1 - \frac{1}{a} \right) - d + 2 \right) \left(n \left(1 - \frac{1}{a} \right) - d + 1 \right) \right]^{\frac{1}{2}} \left(\sum_{i=1}^n x_i^{-a} \right)^{-1}. \quad (24)$$

From equation (10), on using (21), the Bayes estimator of θ under entropy loss function is given by

$$\hat{\theta}_E = \left(n \left(1 - \frac{1}{a} \right) - d \right) \left(\sum_{i=1}^n x_i^{-a} \right)^{-1}. \quad (25)$$

From equation (12), on using (19) and (21), the Bayes estimator of θ under K-loss function is given by

$$\hat{\theta}_K = \left[\left(n \left(1 - \frac{1}{a} \right) - d + 1 \right) \left(n \left(1 - \frac{1}{a} \right) - d \right) \right]^{\frac{1}{2}} \left(\sum_{i=1}^n x_i^{-a} \right)^{-1}. \quad (26)$$

From equation (14), on using (18) and (22), the Bayes estimator of θ under Al-Bayyati's loss function comes out to be

$$\hat{\theta}_{Al} = \left(n \left(1 - \frac{1}{a} \right) - d + c + 1 \right) \left(\sum_{i=1}^n x_i^{-a} \right)^{-1}. \quad (27)$$

5 Posterior Density Under $g_2(\theta)$

Under $g_2(\theta)$, the posterior density of θ , using equation (2), is obtained as

$$f(\theta/\underline{x}) = \frac{\left[\left(\frac{a}{\Gamma\left(1 - \frac{1}{a}\right)} \right)^n \theta^{n\left(1 - \frac{1}{a}\right)} \left(\prod_{i=1}^n x_i^{-a} \right) e^{-\theta \sum_{i=1}^n x_i^{-a}} \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{\alpha-1} e^{-\beta\theta} \right]}{\int_0^\infty \left[\left(\frac{a}{\Gamma\left(1 - \frac{1}{a}\right)} \right)^n \theta^{n\left(1 - \frac{1}{a}\right)} \left(\prod_{i=1}^n x_i^{-a} \right) e^{-\theta \sum_{i=1}^n x_i^{-a}} \frac{\beta^\alpha}{\Gamma(\alpha)} \theta^{\alpha-1} e^{-\beta\theta} \right] d\theta}$$

$$\begin{aligned}
 &= \frac{\theta^{n\left(1-\frac{1}{a}\right)+\alpha-1} e^{-\left(\beta+\sum_{i=1}^n x_i^{-a}\right)\theta}}{\int_0^{\infty} \theta^{n\left(1-\frac{1}{a}\right)+\alpha-1} e^{-\left(\beta+\sum_{i=1}^n x_i^{-a}\right)\theta} d\theta} \\
 &= \frac{\theta^{n\left(1-\frac{1}{a}\right)+\alpha-1} e^{-\left(\beta+\sum_{i=1}^n x_i^{-a}\right)\theta}}{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha\right) \left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{n\left(1-\frac{1}{a}\right)+\alpha}} \\
 &= \frac{\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{n\left(1-\frac{1}{a}\right)+\alpha}}{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha\right)} \theta^{n\left(1-\frac{1}{a}\right)+\alpha-1} e^{-\left(\beta+\sum_{i=1}^n x_i^{-a}\right)\theta} \tag{28}
 \end{aligned}$$

Theorem 2. On using (28), we have

$$E\left(\theta^c\right)=\frac{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha+c\right)}{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha\right)}\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{-c} . \tag{29}$$

Proof. By definition,

$$\begin{aligned}
 E\left(\theta^c\right) &= \int \theta^c f(\theta/\underline{x}) d\theta \\
 &= \frac{\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{n\left(1-\frac{1}{a}\right)+\alpha}}{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha\right)} \int_0^{\infty} \theta^{n\left(1-\frac{1}{a}\right)+\alpha+c-1} e^{-\left(\beta+\sum_{i=1}^n x_i^{-a}\right)\theta} d\theta \\
 &= \frac{\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{n\left(1-\frac{1}{a}\right)+\alpha}}{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha\right)} \times \frac{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha+c\right)}{\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{n\left(1-\frac{1}{a}\right)+\alpha+c}}
 \end{aligned}$$

$$= \frac{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha+c\right)}{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha\right)}\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{-c}.$$

From equation (29),for $c = 1$, we have

$$E(\theta) = \left(n\left(1-\frac{1}{a}\right)+\alpha\right)\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{-1}. \tag{30}$$

From equation (29),for $c = 2$, we have

$$E(\theta^2) = \left[\left(n\left(1-\frac{1}{a}\right)+\alpha+1\right)\left(n\left(1-\frac{1}{a}\right)+\alpha\right)\right]\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{-2}. \tag{31}$$

From equation (29),for $c = -1$, we have

$$E\left(\frac{1}{\theta}\right) = \frac{1}{\left(n\left(1-\frac{1}{a}\right)+\alpha-1\right)}\left(\beta+\sum_{i=1}^n x_i^{-a}\right). \tag{32}$$

From equation (29),for $c = c + 1$, we have

$$E(\theta^{c+1}) = \frac{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha+c+1\right)}{\Gamma\left(n\left(1-\frac{1}{a}\right)+\alpha\right)}\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{-(c+1)}. \tag{33}$$

6 Bayes Estimators Under $g_2(\theta)$

From equation (6), on using (30), the Bayes estimator of θ under squared error loss function is given by

$$\hat{\theta}_S = \left(n\left(1-\frac{1}{a}\right)+\alpha\right)\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{-1}. \tag{34}$$

From equation (8), on using (31), the Bayes estimator of θ under precautionary loss function is obtained as

$$\hat{\theta}_P = \left[\left(n\left(1-\frac{1}{a}\right)+\alpha+1\right)\left(n\left(1-\frac{1}{a}\right)+\alpha\right)\right]^{\frac{1}{2}}\left(\beta+\sum_{i=1}^n x_i^{-a}\right)^{-1}. \tag{35}$$

From equation (10), on using (32), the Bayes estimator of θ under entropy loss function is given by

$$\hat{\theta}_E = \left(n \left(1 - \frac{1}{a} \right) + \alpha + 1 \right) \left(\beta + \sum_{i=1}^n x_i^{-a} \right)^{-1} \quad (36)$$

From equation (12), on using (30) and (32), the Bayes estimator of θ under K-loss function is given by

$$\hat{\theta}_K = \left[\left(n \left(1 - \frac{1}{a} \right) + \alpha \right) \left(n \left(1 - \frac{1}{a} \right) + \alpha - 1 \right) \right]^{\frac{1}{2}} \left(\beta + \sum_{i=1}^n x_i^{-a} \right)^{-1} \quad (37)$$

From equation (14), on using (29) and (33), the Bayes estimator of θ under Al-Bayyati's loss function comes out to be

$$\hat{\theta}_{Al} = \left(n \left(1 - \frac{1}{a} \right) + \alpha + c \right) \left(\beta + \sum_{i=1}^n x_i^{-a} \right)^{-1} \quad (38)$$

7 Conclusion

In this paper, we have obtained a number of estimators of parameter of length-biased weighted Frechet distribution. In equation (4) we have obtained the ML estimator of the parameter. In equations (23), (24), (25), (26) and (27) we have obtained the Bayes estimators under different loss functions using quasi prior. In equations (34), (35), (36), (37) and (38) we have obtained the Bayes estimators under different loss functions using gamma prior. In the above equation, it is clear that the Bayes estimators depend upon the parameters of the prior distribution. We therefore recommend that the estimator's choice lies according to the value of the prior distribution which in turn depends on the situation at hand.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Reyad HM, Hashish AM, Othman SA, Allam SA. The length-biased weighted Frechet distribution: Properties and estimation. *International Journal of Statistics and Applied Mathematics*. 2018;3(1):189-200.
- [2] Zellner A. Bayesian estimation and prediction using asymmetric loss functions. *Jour. Amer. Stat. Assoc.* 1986;91:446-451.
- [3] Basu AP, Ebrahimi N. Bayesian approach to life testing and reliability estimation using asymmetric loss function. *Jour. Stat. Plann. Infer.* 1991;29:21-31.
- [4] Norstrom JG. The use of precautionary loss functions in Risk Analysis. *IEEE Trans. Reliab.* 1996;45(3):400-403.
- [5] Calabria R, Pulcini G. Point estimation under asymmetric loss functions for left truncated exponential samples. *Comm. Statist. Theory & Methods*. 1994;25(3):585-600.
- [6] Dey DK, Ghosh M, Srinivasan C. Simultaneous estimation of parameters under entropy loss. *Jour. Statist. Plan. And infer.* 1987;347-363.

- [7] Dey DK, Pei-San Liao Liu. On comparison of estimators in a generalized life Model. *Microelectron. Reliab.* 1992;32 (1/2):207-221.
- [8] Wasan MT. *Parametric Estimation*. New York: McGraw-Hill; 1970.
- [9] Al-Bayyati. Comparing methods of estimating Weibull failure models using simulation. Ph.D. Thesis, College of Administration and Economics, Baghdad University, Iraq; 2002.

© 2021 Rao and Pandey; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)
<http://www.sdiarticle4.com/review-history/65210>