



Effect of Substituting Poultry Waste Meal for Corn in the Diet of the African Clariid Catfish, *Clarias gariepinus* (Burchell 1822) Juveniles

Bernadine Wuraola Obe^{1*}

¹*Department of Fisheries and Aquaculture Management, Faculty of Agricultural Sciences, Ekiti State University, P.M.B. 5363, Ado-Ekiti, Ekiti State, Nigeria.*

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/JEAI/2019/v40i430372

Editor(s):

(1) Dr. Slawomir Borek, Professor, Department of Plant Physiology, Adam Mickiewicz University, Poland.

Reviewers:

(1) Önder Yildirim, Mugla Sıtkı Kocman University, Turkey.

(2) Byron Baron, University of Malta, Malta.

(3) Mariana Lins Rodrigues, Universidade Estadual do Oeste do Paraná, Brasil.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50694>

Original Research Article

Received 02 June 2019
Accepted 28 August 2019
Published 13 September 2019

ABSTRACT

The objective of this study was to evaluate the effect of poultry waste meal (PWM) on growth of *Clarias gariepinus* juveniles. The growth response of juveniles of average weight 6.5g fed poultry waste meal (PWM) was studied for 70 days. The poultry waste meal comprised the droppings from the layers pen, some maggots, broken eggs and feeds that fell during the course of feeding. The proximate analysis of PWM had crude protein value of 9.795 %, crude fibre 8.700 %, ash 25.174 %, moisture 10.794%, carbohydrate 44.286% and fat 1.250% respectively. Five diets were formulated; which were Diet A (Control: with 0% inclusion level of poultry waste meal, B (25%), C (50%), D (75%), and E with 100% inclusion level (total replacement with poultry waste meal). There were no significant differences ($p>0.05$) between the control and all other diets in the values recorded for growth and nutrient utilization, even up to 100% substitution of maize with PWM. The cost of feed production decreased with increase in inclusion levels of PWM in the diets. The results of the study showed that the use of PWM could be considered in the diet of *C. gariepinus* even up to 100% substitution level; considering the huge cost of maize and competition for its use whereas PWM is obtainable at little or no cost.

*Corresponding author: E-mail: bernadine.obe@eksu.edu.ng;

Keywords: Poultry waste meal; substitution; maize; *Clarias gariepinus*; juveniles.

1. INTRODUCTION

Fish is one of the most highly consumed proteins because of the quality and characteristics of its flesh composition and easy accessibility, therefore it has little or no religious or regional bias [1], therefore the demand for it is on the increase even as the human population is increasing. Capture fisheries were relied upon in the past to meet fish demand but recent trends all over the world points to a decline in landing from capture fisheries which are an indicator that fish stock have approached or even exceeded the point of maximum sustainable yield [2], hence further increase in capture fisheries are not anticipated under the current global condition [3]. With this trend, aquaculture has become the proposed solution to bridge the fish demand-supply gap.

A major aspect of aquaculture is feeding. Fish nutritionists have demonstrated increasing interest in research focused on reducing the cost of the most expensive ingredients by alternative nutrient sources such as replacing a fish meal or other conventional feedstuffs with either plant protein sources or other unconventional feedstuffs [4,5,6]. Maize is one of such expensive and conventional ingredient because there is great competition for its use by both the human food and animal feed industries. For a successful fish culturing project, the optimum dietary requirement at a reduced production cost is essential in order to achieve maximum profit, therefore reducing feed cost is a major challenge in aquaculture nutrition.

Poultry waste meal (PWM) is an agricultural waste and a potential feedstuff which could be a source of protein or energy depending on its composition. [7] observed that agricultural wastes' compositions tend to vary as it will depend on the system and type of agricultural activities from which they are obtained and they can be in the form of liquids, slurries, or solids. [7] further stated that these agricultural wastes are the non-product outputs of production and processing of agricultural products that may contain material that can benefit man or can be collected and processed for beneficial use at a minimal cost. The poultry waste meal used in this study comprised of left overfeeds, broken poultry egg parts and poultry faeces which was found on analysis to have a proximate composition similar to the yellow maize.

The African Clariid catfish, *Clarias gariepinus* (Burchell, 1822) is one of the most popularly cultured fish species in Nigeria because of its many aquaculture potentials [8,9]. Information on the effect of poultry waste meal, when incorporated into fish diets, is scarce.

This study, therefore, is focused on the use of poultry waste meal as a replacement for maize, in the diet of African catfish *Clarias gariepinus* juveniles so as to reduce feed cost.

2. MATERIALS AND METHODS

The study was carried out at the Department of Fisheries and Aquaculture Management, Ekiti State University Ado-Ekiti, Nigeria Research Laboratory.

The dietary ingredients: Fish meal, soybean meal, vitamin premix, bone meal and starch were purchased from Metrovet Veterinary Shop, G.R.A. Ado Ekiti and Mercy Agricultural Business Services, Ado Ekiti.

2.1 Collection and Preparation of Poultry Waste Meal

The poultry waste was collected from Ekiti State University Poultry Farm. The poultry waste comprised of poultry droppings of layers, some maggots, some broken eggs and feeds that fell during the course of feeding. The poultry waste was sun-dried at a temperature of 28°C for three days and then ground to flour using the grinding machine. The ground poultry waste was then taken to the laboratory for proximate composition before being incorporated into the experimental diets.

2.2 Preparation of Experimental Diets

The dietary ingredients were measured as contained in Table 1 with poultry waste meal (PWM) substituted for maize at inclusion levels 0%, 25%, 50%, 75% and 100% for diet A (control), B, C, D and E respectively. Starch was added to act as a binder and it was pelletized with a locally fabricated pelleting machine of 3mm die size. The pellets were sun-dried and packed in well-labelled airtight containers and stored in a cool and dry place.

2.3 Procurement of Fish

One hundred and fifty juvenile *Clarias gariepinus* of the average weight of 6.5 g were purchased

from Federal Ministry of Agriculture GRA, Ado Ekiti. The fish were starved for 24 hours and allowed to acclimate to the new environment after which a Coppens feed was used to feed the fish for three days prior to the beginning of the actual feeding experiment.

Ten fish were randomly stocked per aquarium in triplicate groups for each treatment and fed twice daily to satiation at 8.00 to 9.00 and 17.00 to 18.00 hours for 70 days. Weighing of fish in each aquarium was carried out in batches every fortnight and feed was adjusted to accommodate the increase in body weights of the fish. Faeces and feed remnants were siphoned out every day to prevent fouling.

2.4 Determination of Growth Performance of Test Fish

Growth performance was determined as follows [10]:

- i Weight gain = final weight of fish (W_2)- Initial weight (W_1)
- ii Specific growth rate (SGR) = Log_e final weight- Log_e initial weight \times 10 Rearing period (Days)
- iii Protein efficiency ratio (PER) = fish weight gain (g) Protein consumed (g)
- iv Feed conversion ratio (FCR) = weight of feed (g) Fish weight gain (g)

2.5 Proximate Analyses

The proximate analyses of poultry waste meal (PWM) and that of the fish after the experiment were determined using the method of [11]. Parameters determined were: moisture content,

crude protein, lipid, ash, crude fibre and NFE (carbohydrate).

2.6 Cost of Experimental Diets

The cost of producing 1 kg of the different feeds with PWM at different inclusion levels was calculated and compared with the production cost of 1 kg of the control diet which has no PWM. Costing was done according to the prevailing market prices of ingredients that were used in diets at the time of the experiment.

2.7 Statistical Analysis

Data on growth parameters were subjected to one - way analysis of variance (ANOVA) to test for the significant difference in the means while means which were significantly different were separated using Duncan's multiple range test. The analysis was performed using the SPSS (Statistical Package for Social Sciences) version 21. Significant level was set at $p \geq 0.05$.

3. RESULTS

3.1 Proximate Composition of Poultry Waste Meal

The result of the proximate composition of poultry waste meal is shown in Table 2.

3.2 Growth and Nutrient Utilization of the Experimental Fish

Table 3 shows the growth and nutrient utilization of the experimental fish. There were no significant differences ($p > 0.05$) between all variables of growth when using the diets containing PWM and control.

Table 1. Gross composition of poultry waste meal (PWM) diets (g/100 g)

Levels of inclusion	0% PWM	25% PWM	50% PWM	75% PWM	100% PWM
Fishmeal (65.5%)	32.00	32.00	32.00	32.00	32.00
SBM (45%)	31.5	31.5	31.5	31.5	31.5
Yellow maize (10%)	28.5	22.8	17.1	11.4	5.7
PWM (9.8%)	-	5.7	11.4	17.1	22.8
Veg. oil	2.5	2.5	2.5	2.5	2.5
Bone meal	1.00	1.00	1.00	1.00	1.00
*VIT. Premix	2.00	2.00	2.00	2.00	2.00
Starch	2.5	2.5	2.5	2.5	2.5

*Each kg contains: Vit. A: 4,000,000IU; Vit. B: 800,000IU; Vit. E: 16,000 mg; Vit. K3: 800 mg; Vit. B1: 600 mg; Vit. B2: 2,000 mg; Vit. B6: 1,600 mg; Vit. B12: 8 mg; Niacin: 16,000 mg; Caplan: 4,000 mg; Folic Acid: 400 mg; Biotin: 40 mg; Antioxidant: 40,000 mg; Chlorine chloride: 120,000 mg; Manganese: 32,000 mg; Iron: 16,000 mg; Zinc: 24,000 mg; Copper: 32,000 mg; Iodine: 320mg; Cobalt: 120 mg; Selenium: 800 mg manufactured by DSM Nutritional products Europe Limited, Basle, Switzerland.

Where: SBM = Soybean meal, PWM = Poultry waste meal

Table 2. Proximate composition of poultry waste meal (%) dry weight

FAT	1.250
Crude Fibre	8.700
Protein	9.795
Moisture	10.794
Ash	25.174
Carbohydrate	44.286

Laboratory analysis, 2019

3.3 Carcass Composition of the Experimental Fish

The carcass composition of the experimental fish is given in Table 4. Fish fed with diet containing 25% PWM had the highest crude protein and ash content value. Fish fed 100% PWM presented higher moisture values than those fed with the control diet. Ether extract and NFE were higher in control diet-fed fish.

3.4 Water Parameters during the Experimental Period

The temperature and the dissolved oxygen throughout the period of the experiment ranged from 26.120-26.272°C and 6.266-7.300 mg/litre respectively while pH values ranged between 6.300-6.733.

3.5 Cost of Experimental Diet

The cost of replacing maize in the diet of *Clarias gariepinus* is shown in Table 5. The cost of the diet decreased with increasing level of inclusion of poultry waste meal.

Cost analysis of replacing maize with poultry waste meal showed that the cost of production of 1kg feed reduced as the level of inclusion of poultry waste meal increased.

4. DISCUSSION

The experimental fish species (*Clarias gariepinus*) readily accepted the experimental diets and showed evidence of good growth as attested to by the final weight and weight gain recorded in the experiment. This is an indication that poultry waste meal could be nutritious and well utilized for conversion to flesh in the diets of fish. This is in agreement with the work of [7] who reported that agricultural wastes may contain materials that can be collected and processed for beneficial use at a minimal cost. It is also a common practice in integrated poultry-cum- fish

farming that poultry wastes could serve as the main source of food for the fish all through the culture period, bringing about adequate growth at little cost with no adverse effect on the fish health. The result of this study further corroborates the work of [12] who reported that some industrial wastes of the food industry (biscuit, Indomine® and Gala® wastes) had equal potentials in replacing maize as an energy supplement when incorporated to the *Clarias gariepinus* feed with positive effects on the growth and haematological parameters of the fish.

The result of the proximate analysis of PWM in this study showed that it had crude protein level and other nutrients very similar to that of yellow maize. This is an indication that it could conveniently replace maize in this experiment with no adverse effects on the fish growth as attested to by the result of this study. In nutrition experiments, usually feed ingredients with comparable nutrients and energy compositions are used for substituting the other so that there will be no deficiency symptoms manifesting in the animal being fed as a result of the substituted ingredient or introduce bias in the results obtained in the experiment. The result of proximate composition of PWM obtained from laboratory analysis in this study however is in contrast to the work of [13] who reported a crude protein value of 28.6% on analysis of poultry waste. The differences in values could be due to collection and processing methods. [7] also reported that agricultural wastes usually tend to vary in composition depending on the constituents.

From the growth and nutrient utilization results obtained in the study, there were no significant differences ($p>0.05$) between the results obtained in the control and all the other diets. Considering the high cost of maize in the market and the fact that PMW could be obtained at little or no cost, its incorporation could be a way of converting waste to wealth. Incorporating it in the diet of *Clarias gariepinus* would greatly reduce the cost of feed and consequently the cost of production and increase the profit margin of farmers. Several authors have also worked on the replacement of maize with lesser-used ingredients with varying levels of success [14, 12,6,15,16].

The experimental water condition was maintained at temperature values between 26.12 and 26.27°C, dissolved oxygen values between

6.26 and 7.30 mg/l and pH values between 6.3 and 6.7 5. These values were within the standard values recommended for warm water fish culture and supported good fish production [17,18].

The result of the carcass analysis also showed that the crude protein for all the fish fed the experimental diets were higher than that of the control which showed that they retained the protein in their carcass than the fish fed the maize-based diets.

The cost of replacing maize with poultry waste meal shows that the cost of production of 1 kg feed reduced as the level of inclusion of poultry waste meal increased. Profitability and viability of a fish farming enterprise depend largely on the total cost of fish feeds as feeding cost represents

the most expensive component of the fish farming enterprise, therefore, the more the inclusion level of PWM in this experiment, the more profitable and viable the project would be. This is in line with the findings of [12] who reported that the use of some industrial wastes in replacing maize in the diet of *Clarias gariepinus* all reduced total feed cost per kg by at least 30%. The fish farming sector is currently faced with the challenge of the inadequate and prohibitive cost of quality fish feeds, therefore, it is pertinent that the use of alternative sources of nutrients that ordinarily pass as waste and usually discarded such as PWM used in this experiment be explored as it is not competed for like maize. These agricultural wastes could be procured at little or no cost as they are categorized as waste products meant to be discarded.

Table 3. Growth and nutrient utilization of *Clarias gariepinus* fed with poultry waste meal

	Control	25% PWM	50% PWM	75%PWM	100% PWM
Mean initial weight	6.553±0.006 ^a	6.557±0.006 ^a	6.550±0.000 ^a	6.557±0.006 ^a	6.553±0.006 ^a
Mean final weight	15.780±2.130 ^a	14.940±0.949 ^a	13.597±1.346 ^a	14.477±1.476 ^a	13.610±0.582 ^a
Mean weight gain	9.227±2.129 ^a	8.383±0.947 ^d	7.047±1.346 ^a	7.920±1.475 ^a	7.057±0.583 ^a
Average daily weight gain	0.1318±0.031 ^a	0.1198±0.017 ^a	0.1007±0.006 ^a	0.1131±0.025 ^a	0.1008±0.010 ^a
Specific growth rate (SGR)	1.247±0.203 ^a	1.183±0.102 ^a	1.040±0.148 ^a	1.127±0.146 ^a	1.047±0.060 ^a
Protein efficiency ratio (PER)	0.692±0.169 ^a	0.630±0.0701 ^a	0.529±0.101 ^a	0.594±0.111 ^a	0.529±0.044 ^a
Feed conversion ratio (FCR)	1.880±4.814 ^a	2.003±0.215 ^a	2.430±0.521 ^a	2.152±0.393 ^a	2.373±0.193 ^a

Means and standard deviation along the same column followed by the same superscripts are not significantly different ($p>0.05$)

Table 4. Carcass composition of experimental fish (% Dry weight)

	Control	25% PWM	50% PWM	75%PWM	100% PWM
Moisture	6.828±0.047 ^b	5.890±0.045 ^a	6.607±0.025 ^{ab}	6.829±0.104 ^b	7.150±0.754 ^b
Crude protein	59.195±1.318 ^a	70.172±0.962 ^c	64.083±0.818 ^b	64.998±0.123 ^b	62.098±2.341 ^{ab}
Ether extract	15.248±0.543 ^c	12.260±0.514 ^a	14.686±0.675 ^{bc}	13.234±0.320 ^{ab}	14.067±1.115 ^{abc}
Ash	4.998±0.064 ^b	5.291±0.004 ^c	4.990±0.009 ^b	4.856±0.009 ^a	5.059±0.075 ^b
NFE	13.730±0.756 ^d	6.411±0.442 ^a	9.684±0.136 ^b	10.082±0.348 ^b	11.619±0.403 ^c

Means and standard deviation along the same column followed by the same superscripts are not significantly different ($p>0.05$)




Fig. 1. Graph of the weight increase of experimental fish

Table 5. Cost of experimental diet

Ingrediens	control(N)	25%PWM(N)	50%PWM(N)	75%PWM(N)	100%PWM(N)
Fish meal	1080	1080	1080	1080	1080
Poultry waste meal	-	7.125	14.25	21.38	28.6
Soybean meal	143.33	143.33	143.33	143.33	143.33
Yellow maize	79.8	63.84	47.88	31.92	15.96
Vegetable oil	25.00	25.00	25.00	25.00	25.00
Bone meal	1.5	1.5	1.5	1.5	1.5
Vitamin premix	82.5	82.5	82.5	82.5	82.5
Starch	6.25	6.25	6.25	6.25	6.25
Total	1418.41	1409.55	1400.71	1391.88	1383.14

5. CONCLUSION

Considering the high cost of maize in both local and international markets and the competition for its use in both human foods and livestock feeds, the use of PMW is a good alternative for reducing the cost of production and making fish farming more profitable. It will also serve as a way of converting waste to wealth.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

- Eyo AA. Fish Processing Technology in the Tropics. University of Ibadan Press, Ibadan, Nigeria; 2001.
- Faturoti EO. Fisheries potential and investment opportunities in Nigeria. Paper presented at the Fisheries Society of Nigeria (FISON) Investors Workshop held at NIOMR Lagos. 1999;22.
- Ounham RA, Majumdar K, Hallerman E, Bartly D, Mair G, Hulata G, Liu Z, Pongthan N, Bakos J, Penman D, Gupta M, Rothlisberg P, Hoerstgen-Schwark G. Review of the status of aquaculture genetics in the third millenium. Technical Proceedings of the Conference on Aquaculture, Bangkok, Thailand. 2001; 137-166.
- Fagbenro OA, Balogun AM, Ibronke AA, Fasina FA. Nutritional value of some amphibian meals in diets for *Clarias gariepinus* (Burchell 1822) (Siluriformes: Clariidae). Journal of Aquaculture in the Tropics. 1993;8:95-101.

5. Adeparusi EO, Olute BW. Effect of methionine supplemented toasted lima bean (*Phaseolus lunatus*) diets on the Growth of *Oreochromis niloticus*. Applied Tropical Agriculture. 2001;5(2):113-117.
6. Obe BW. Growth performance and nutrient utilization of catfish hybrid (*Heterobranchus bidorsalis* X *Clarias gariepinus*) fed fermented sorghum (*Sorghum bicolor*) waste meal diets. International Journal of Applied Science and Technology. 2014;4(3):130-136
7. Obi FO, Ugwuishiwu BO, Nwakaire JN. Agricultural Waste Concept, Generation, Utilization and Management. Nigerian Journal of Technology (NIJOTECH) Faculty of Engineering, University of Nigeria, Nsukka. 2016;35(4):957-964.
8. Fagbenro OA, Nwanna LC, Adeparusi EO, Adebayo OT, Fapohunda OO. An overview of animal feed industry and dietary substitution of feedstuffs for farmed fish in Nigeria. In: Crops: Growth, quality and biotechnology (current status and future prospects) (Ramdane Dris, editor). WFL Publisher, Helsinki, Finland. 2005;91-107.
9. Fasakin EA, Davies SJ, Akegbejo-Samsons Y. Efficacy of maize gluten supplemented with crystalline l-lysine in the diets for the African clariid catfish *Clarias gariepinus* (Burchell,1822). Journal of Animal and Veterinary Advances. 2006; 5(3):237-243.
10. Steffens W. Principles of fish nutrition. Ellis Horwood Ltd. England; 1989.
11. A.O.A.C. Association of Official Analytical Chemists. Official methods of analysis, 15th Edition. Washington D.C. United States of America. 1990;1230.
12. Agbebi OT, Ilesanmi AI, Alegbeleye WO, Odulate DO, Obasa SO, Olaoye OJ. Growth and haematological effects of replacing maize with feed wastes on the Juvenile *Clarias gariepinus* (Burchell, 1822). Journal of Fisheries and Aquatic Science, 2013;8:535-543.
13. Obasa SO, Alegbeleye WO, Amole JB. Dried poultry manure meal as a substitute for soybean meal in the diets of African Catfish (*Clarias gariepinus*) (Burchell 1822) Advanced Fry. Turkish Journal of Fisheries and Aquatic Sciences. 2009;9:121-124.
14. Obasa SO, Alegbeleye WO, Akinyemi AA, Idowu AA, Bamidele NA, Adesanya AN. Replacement of maize meal by toasted African breadfruit (*Treculia africana*) seed meal in the diet of *Clarias gariepinus* (Burchell 1822) Fingerlings. Livestock Research for Rural Development LRRD Newsletter. 2013;25(6).
15. Fakunmoju O, Babalola A, Ijimakinde B, Anjola OA, Orowole PF. Effect of substituting maize with bambara (*Voandzeia subteranea* Thouars) waste meal in the practical diets of *Tilapia niloticus* fingerlings. Journal of Fisheries and Aquatic Science. 2016;11: 185-189.
16. Obe BW. Use of fermented wheat (*Triticum spp*) waste meal in the diet of *Clarias gariepinus* Fingerlings. American Academic and Scholarly Research Journal. 2017;9(4):91-100.
17. Swann L. A basic overview of aquaculture: History, water quality, types of aquaculture, and production methods. Illinois-Indiana Sea Grant Program Extension Bulletin AS-457 and IL-IN-SG-E-90G:\Final 13-9-19\JEAI_c++\GP-4-50694-PH---\Final publication-2. Purdue University. West Lafayette, Indiana; 1990.
18. Landau M. Introduction to Aquaculture. John Wiley and Sons Inc. New York; 1992.

© 2019 Obe; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50694>