



# Exact Solution of Particle's Occupation Number in the Lambert Boltzmann Distribution using Lambert W Function

Chokri Hadj Belgacem<sup>1,2\*</sup>

<sup>1</sup>Department of Physics, Science College at Yanbu, Taibah University, KSA.

<sup>2</sup>Department of Physics, LMCN, Faculty of Science, Monastir University, Tunisia.

## Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

## Article Information

### Editor(s):

(1) Dr. Yong X. Gan, Professor, Department of Mechanical Engineering, California State Polytechnic University, Pomona, USA.

### Reviewers:

(1) Abdullah Sonmezoglu, Yozgat Bozok University, Turkey.

(2) Adel H. Phillips, Ain-Shams University, Egypt.

(3) Snehadri B. Ota, Institute of Physics, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48382>

Received 31 January 2019

Accepted 17 April 2019

Published 25 April 2019

Original Research Article

## ABSTRACT

By applying the exact Stirling formula and the exact function  $\ln(n!)$ , the occupation number of particles was calculated based on the statistics of Boltzmann distribution for a finite number of particles  $n$ . The exact analytical expression of occupation number of particles ( $n_i$ ) is found in terms of the Lambert W function and is more general than that usually calculated by the standard Boltzmann distribution based on the Stirling approximation  $\ln(n!) = n \ln(n) - n$ . The new expression in the exact and algebraic closed form eliminates the need for the complex iterative computation. Its high accuracy is proved by a comparison of calculating occupation number of particles ( $n_i$ ) with respective numerical solution.

**Keywords:** Occupation number of particles; Lambert Boltzmann distribution; finite number of particles; Lambert W function.

\*Corresponding author: Email: Chokri.hadjbelgacem@laposte.net, chokri.hadbelgacemj@laposte.net;

### 1. INTRODUCTION

The Stirling approximation to  $\ln(n!)$  is typically introduced to physical chemistry students as a step in the derivation of the exponential Boltzmann distribution. However, naïve application of this approximation leads to incorrect conclusions. For example student of physical chemistry are often introduced to the statistical treatment of the occupation number  $n_i$  as given by Boltzmann's distribution equation:

$$n_i = \exp[-\alpha - \beta \varepsilon_i] \tag{1}$$

Using Stirling's approximation [1,2]

$$\ln(n!) = n \ln(n) - n \tag{2}$$

The applicability range of Eq. (2) is limited to very large values of  $n$ , the exponential Boltzmann distribution is limited strictly to  $n \gg 1$  and applies only for  $\varepsilon$  not too large. However, the distribution is used in the literature even for  $\varepsilon \rightarrow \infty$  and  $n \rightarrow 0$ , where Eq. (1) does not be appropriate. It is interesting that Stirling's approximation, Eq.(1), fails and the more precise Stirling formula, Eq.(3) [2] is required to determine the occupation number  $n_i$ .

$$\ln(n!) = \left(n + \frac{1}{2}\right) \ln(n) - n + \frac{1}{2} \ln(2\pi) \tag{3}$$

For a finite number of particles, the occupation number of particle has been determined by Kakorin [3], but no details, analysis or more explanation and discussion were made for how determining it.

In this paper, basing on the work of Kakorin [3], we present more details, analysis and discussion to calculate the expression of the occupation number  $n_i$  of particles based on Boltzmann statistics for a finite number of particles. We compare the obtained result with this obtained by usual exponential Boltzmann statistics using the Stirling approximation  $\ln(n!) = n \ln(n) - n$ .

### 2. METHODOLOGY

By applying the exact Stirling formula and the exact function  $\ln(n!)$  based on the statistics of Boltzmann distribution for a finite number of particles  $n$ , we calculate the occupation number

of particles. Analytical expression of  $n_i$  was expressed in terms of the primary branch of the Lambert function  $W_0$ .

Using numerical method as Newton–Raphson's method, the occupation number of particles was also calculated and compared with this obtained by applying exact Stirling formula.

### 3. RESULTS AND DISCUSSION

Based on the Boltzmann statistics with a finite number of particles  $n$ , the expression of occupation number  $n_i$  is [3]

$$\frac{1}{2n_i^*} = -\ln(n_i^*) - A \tag{4}$$

where  $A$ , represent some constant in the Boltzmann statistics.

Eq. (4) can be solved numerically by an iterative method. This is not necessary; the exact solution of this equation is given by the so-called Lambert W-function [4]. This function was postulated to solve the equation:

$$W(z) \exp W(z) = z \tag{5}$$

The Lambert W-function allows the explicit solution of entire classes of differential equations, which actually only could be solved numerically and is experiencing today a renaissance in various fields of sciences and engineering [5-12].

The expression of the occupation number of particles is given by:

$$\frac{1}{2n_i} = -\ln(n_i) - \alpha - \beta \varepsilon_i \tag{6}$$

Where  $\varepsilon_i$  is the energy of state (i),  $\beta = 1/kT$  ( $k$  is the Boltzmann constant and  $T$  is the temperature and  $\alpha$  is the Lagrange multiplier).

Eq. (6) is a transcendental equation that can be solved exactly with the results written in closed form in terms of the Lambert W function using the approach proposed by Hadj Belgacem [13-17].

For further calculation, we introduce the abbreviations:

$$A = \alpha + \beta \varepsilon_i \tag{7} \quad \text{Exp}(W(z) + \ln(W(z)) - \ln(z)) = 1 \tag{13}$$

and  $z = -\frac{\text{Exp}(A)}{2}$  (8)  $W(z) \text{Exp}W(z) = z$  (14)

We obtain the implicit equation

$$\frac{1}{2n_i} = -\ln(n_i) - A \tag{9}$$

For solving Eq. (9) we suppose that

$$n_i = \text{Exp}(-A) \frac{z}{W(z)} \tag{10} \quad n_i = -\frac{1}{2W\left(-\frac{1}{2}\text{Exp}(\alpha + \beta \varepsilon_i)\right)} \tag{15}$$

Inserting Eq. (10) into Eq. (9) yields

$$\begin{aligned} \frac{1}{2 \text{Exp}(-A) \frac{z}{W(z)}} &= -\ln\left(\text{Exp}(-A) \left(\frac{z}{W(z)}\right)\right) - A \\ \frac{W(z)}{2 \text{Exp}(-A) \left(-\frac{\text{Exp}(A)}{2}\right)} &= -\ln(\text{Exp}(-A)) - \ln\left(\frac{z}{W(z)}\right) - A \\ W(z) + \ln\left(\frac{W(z)}{z}\right) &= 0 \end{aligned} \tag{11}$$

Rearranging Eq. (11) and employing an exponential in Eq. (13)

$$W(z) + \ln(W(z)) - \ln(z) = 0 \tag{12}$$

We find that Eq. (14) represents the definition of the Lambert W-function, as already established in Eq. (5).

Consequently, our supposition in Eq. (10) is justified.

The exact solution for the occupation number  $n_i$  is after resubstitution for  $z$  and replacing  $A$  is

The Lambert W-function is a complex and multi-valued function with an infinite number of branches, only two of them having real values. If  $x$  is real, then for  $-\frac{1}{e} \leq x \leq 0$ , there are two possible real values of  $W(x)$ , as displayed in reference [4]. The branch satisfying  $-1 \leq W(x)$  is denoted  $W_0(x)$ ; the branch satisfying  $W(x) \leq -1$  is denoted  $W_{-1}(x)$ . Where are  $W(-\frac{1}{e}) = -1$  and  $W(0) = 0$ . Both real branches  $W_0(x)$  and  $W_{-1}(x)$ , for  $x$  real are presented in Fig. 1.






Fig. 1. The two branches of the Lambert W function  $W_{-1}(x)$  in blue color and  $W_0(x)$  in red



**Fig. 2. (a) Exponential Boltzmann distribution  $n_i$  from Eq1 versus  $\beta\epsilon_i$ , where  $\epsilon_i$  is the energy of the state and  $\beta = \frac{1}{kT}$ ; (b) Lambert –Boltzmann distribution  $n_i$  Eq.14 versus  $\beta\epsilon$  with the primary branch of the Lambert W function  $W_0$  and (c) Lambert–Boltzmann distribution  $n_i$  Eq.14 versus  $\beta\epsilon_i$  calculated with the secondary branch of the Lambert W function  $W_{-1}$**



**Fig. 3. The occupation number  $n_i$  as a function of  $\beta\epsilon_i$ . The dashed curve gives the occupation number as calculated from numerical solution. The solid curve gives the present result**

For practical application, to find the branch of  $W(x)$  that correctly describes the evolution of the occupation number  $n_i$  additional reasonable considerations are required. At the limit when the occupation number  $n_i$  tend to infinity ( $n_i \rightarrow \infty$ ), the expression of the general occupation number calculated in this work using the exact striling formula must be equal to the usual one calculating by using the striling approximation.

Inserting appropriate parameters  $n_0 = 1000, \alpha = -\ln(n_0) - \frac{1}{2n_0}$  in Eq.(14);

$n_0 = 1000, \alpha = -\ln(n_0)$  in Eq.(1). The occupation number can be represented in Fig. 2 as a function of  $\beta\epsilon_i$ , where  $\epsilon_i$  is the energy of the state and  $\beta = \frac{1}{kT}$ .

To illustrate what kinds of the Lambert W function branch's that computes the real value of  $n_i$ . We compare in the Fig. 2 the evolution of the general occupation number  $n_i$  calculate with the primary branch  $W_0$  (Fig. 2.a) and with the second branch  $W_{-1}$  (Fig. 2.b) using Eq.14 with usual one calculating by using the striling approximation using Eq.(1) (Fig. 2.c).

The comparison of the general occupation number evolution showed in the Figs. (2.a) and

(2.b) with the usual occupation number Fig. (2.c) demonstrate that  $W_0(x)$  is the branch that appropriately describes the evolution of the general occupation number as a function of  $\beta\epsilon_i$ .

To validate the obtained general analytical solution of the occupation number, we compare it in Fig. 3 with the numerical solution of Eq. (3) obtained using the Newton–Raphson’s method. It is easily seen that an excellent agreement is achieved for all values of the  $\beta\epsilon_i$ .

#### 4. CONCLUSION

In summary, the Lambert W-function was successfully used to determine an exact analytical solution for calculating the occupation number  $n_i$  of particles in the case of Boltzmann statistics for a finite number of particles. The exact solution for the occupation number  $n_i$  was expressed in terms of the primary branch of the Lambert function  $W_0$ . Comparing with numerical result shows that the proposed solution is in a good agreement.

Practically, this occupation number  $n_i$  is simple to compute since the lambert w-function is readily available in standard computational packages and can be easily implemented in other mathematical formulas, for example to calculate the partition function  $z$  and the constant volume heat capacity  $C_v$  in boltzmann statistics.

## COMPETING INTERESTS

Author has declared that no competing interests exist.

## REFERENCES

1. Russel DK. The Boltzmann distribution. *Journal of Chemical Education*. 1996;73:299.
2. Wallner AS, Brandt KA. The validity of Stirling's approximation: A physical chemistry project. *Journal of Chemical Education*. 1999;76:1395.
3. Kakorin S, Kakorin S. Revision of Boltzmann statistics for a finite number of particles. *American Journal of Physics*. 2009;77:1.
4. Corless RM, Gonnet GH, Hare DEG, Jeffrey DJ, Kunth DE. On the Lambert W function. *Advances in Computational Mathematics*. 1996;5:329.
5. Williams BW. Exact solutions of a Schrödinger equation based on the Lambert function. *Physics Letters A*. 2005;334:117.
6. Chen R, Zheng X, Deng W, Wu Z. A physics-based analytical solution to the surface potential of polysilicon thin film transistors using the Lambert W function. *Solid-State Electronics*. 2007;51:975.
7. Bian W, He J, Zhang L, Zhang J, Chan M. Sub-threshold behavior of long channel undoped cylindrical surrounding-gate MOSFETs. *Microelectronics Reliability*. 2009;49:897.
8. Morales DA. Exact expressions for the range and the optimal angle of a projectile with linear drag. *Canadian Journal of Physics*. 2005;83:67.
9. Vial A. Horizontal distance travelled by a mobile experiencing a quadratic drag force: Normalized distance and parametrization. *European Journal of Physics*. 2007;28:657.
10. Warburton RDH, Wang J. Analysis of asymptotic projectile motion with air resistance using the Lambert W function. *American Journal of Physics*. 2004;72:1404.
11. Parker GW. Projectile motion with air resistance quadratic in the speed. *American Journal of Physics*. 1977;45:606.
12. Hadj Belgacem C. Determination of commercial silicon diode (40071N) parameters from the I-V and PV characteristics. *Silicon*. 2018;10(4):1469-1474.
13. Hadj Belgacem C. Exact analytical solution for the critical layer thickness of a lattice-mismatched heteroepitaxial layer. *Journal of Electronic Materials*. 2010;39:2248-2250.
14. Hadj Belgacem C, Fnaiech M. Solution for the critical thickness models of dislocation generation in epitaxial thin films using the Lambert W function. *Journal of Materials Science*. 2011;46:1913-1915.
15. Hadj Belgacem C. Range and flight time of quadratic resisted projectile motion using the Lambert W function. *European Journal of Physics*. 2014;35:055025.
16. Hadj Belgacem C. Explicit solution for critical thickness of semicircular misfit dislocation loops in strained semiconductors heterostructures. *Silicon*. 2016;8(3):397-399.
17. Hadj Belgacem C. Analysis of projectile motion with quadratic air resistance from a nonzero height using the Lambert W function. *Journal of Taibah University for Science*. 2017;11:328-331.

© 2019 Belgacem; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/48382>