



Growth and Biomass Distribution in Cocoa Varieties under Different Nitrogen Regimes

Nayana H ^{a*}, Ravi Bhat ^b, Bhavishya ^c, Maruthi Prasad B N ^a,
S Elain Apshara ^c, S V Ramesh ^b and J Venkatesh ^a

^a Department of PSMAC, College of Horticulture, Bengaluru - 560065, India.

^b ICAR-Central Plantation Crops Research Institute, Kasaragod - 671 124, India.

^c ICAR-Central Plantation Crops Research Institute, Regional Station, Vittal - 574 243, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i82729>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/120306>

Original Research Article

Received: 27/05/2024

Accepted: 30/07/2024

Published: 31/07/2024

ABSTRACT

Among the 17 vital plant nutrients, nitrogen (N) is crucial in crop production. It extensively enhances the growth and biomass production of plants by improving photosynthesis and protein synthesis. To study the impact of three different N levels (40, 80, and 120 ppm) on the growth and biomass production of eight cocoa varieties, a hydroponic experiment was carried out at ICAR-CPCRI, Regional Station, Vittal. The results showed that increasing N concentration significantly boosted growth and biomass production. Among the different varieties, VTLCH-1 exhibited the highest plant height, maximum number of leaves, greater stem diameter, and the highest total fresh biomass with the application of 120 ppm nitrogen. Biomass partitioning analysis revealed that higher N levels

*Corresponding author: E-mail: nayanaN32@gmail.com;

increased leaf and stem biomass while reducing root biomass, leading to a higher shoot-to-root ratio. This indicates that N application influences the biomass production and distribution in the plant.

Keywords: Cocoa; nitrogen; growth; biomass partitioning.

1. INTRODUCTION

Cocoa (*Theobroma cacao* L.), native to the Amazon region, is a perennial plant of significant economic importance and one of the most widely traded agricultural commodities globally [1]. As with most crops, cocoa requires nitrogen (N) in the largest quantities [2]. Nitrogen, often referred to as the “life element,” is essential in the composition of various metabolic and active substances in plants. Nitrogen deficiency can lead to weakened root growth, altered root structure, reduced plant biomass, and decreased photosynthesis [3].

Nitrogen is the most limiting nutrient for crop production in many of the tropical soils, and its efficient use is important for the economic sustainability of cropping systems [4]. Nitrogen additions stimulate the growth of young cocoa seedlings and jorquette formation, particularly in interaction with potassium (K) and magnesium (Mg) applications, thereby promoting early closure of the cocoa canopy [5]. High yield in cocoa is contingent upon optimal nitrogen levels, as N supports the vegetative phase, determining the number of cocoa pods produced during the generative stage [6]. Moreover, N is crucial for developing the cocoa canopy and frame prior to pod production, thus enhancing overall growth [7].

Previous studies, such as those conducted by Rieiro et al., have reported significant genotypic differences in the growth and biomass production of cocoa. Understanding these genotypic responses is crucial for optimizing nitrogen management strategies to enhance cocoa productivity and sustainability. The current study aims to address this gap by investigating the effect of different nitrogen levels on the growth, biomass production, and partitioning in cocoa seedlings. By examining these parameters at the eight-month stage, this research seeks to elucidate the interactions between nitrogen availability and genotypic variation in cocoa. The results of this study could provide valuable insights into developing genotype-specific nitrogen management practices, ultimately contributing to more efficient nutrient use and

improved cocoa yields. Despite the recognized importance of N, the varietal differences in response to varying nitrogen doses remain underexplored. Therefore, this study aims to analyze the growth and biomass partitioning in different cocoa varieties subjected to three different nitrogen levels, providing valuable insights into optimizing nitrogen use for improved cocoa cultivation.

2. MATERIALS AND METHODS

The experiment was conducted at the ICAR-Central Plantation Crops Research Institute (CPCRI), Regional Station, Vittal, during 2021-2022, in a glasshouse. The methodology followed was the same as described in an earlier study by Nayana et al. [8], with the only difference being the nitrogen levels. At eight months, growth parameters such as plant height, number of leaves, and stem diameter were recorded. Subsequently, plants were removed from the nutrient solution, and the fresh weights of leaves, stems, and roots were measured. The plant parts were dried in a hot air oven at 65 °C until a constant weight was obtained. Biomass partitioning was determined by measuring the dry weight of different plant parts (roots, stems, leaves) and expressing each as a percentage of the total plant dry weight.

2.1 Statistical Analysis

Data were subjected to analysis of variance (ANOVA) in a factorial design, considering two sources of variation: cocoa varieties and N levels. The significance of differences among the mean values was assessed at 5% significance level.

3. RESULTS

3.1 Growth Parameters

The growth parameters, viz. plant height, number of leaves, and stem diameter, varied significantly among different cocoa varieties, N levels, and their interactions (Table 1). Nitrogen was found to have a profound effect on the growth of cocoa

Table 1. Growth parameters of different cocoa varieties as influenced by varied N levels

Varieties	Plant height (cm)				Number of leaves				Stem diameter (mm)			
	N1	N2	N3	Mean A	N1	N2	N3	Mean A	N1	N2	N3	Mean A
V1	93.00	108.40	125.07	108.82	43.33	54.00	69.33	55.55	10.60	9.67	11.13	10.47
V2	70.57	84.10	76.87	77.18	43.00	44.67	42.00	43.22	9.60	9.43	10.53	9.85
V3	72.00	79.83	91.87	81.23	39.33	44.33	48.67	44.11	8.87	10.87	10.83	10.19
V4	65.43	96.50	102.43	88.12	39.67	47.00	55.00	47.22	8.77	9.87	11.20	9.95
V5	56.33	80.17	97.27	77.92	50.33	42.67	58.00	50.33	9.47	9.60	12.37	10.48
V6	86.67	87.00	98.83	90.83	49.33	44.67	53.67	49.22	10.13	9.80	10.83	10.25
V7	105.00	110.07	85.67	100.25	50.00	63.00	48.67	53.89	10.60	9.70	9.73	10.01
V8	66.90	81.97	96.83	81.90	36.33	37.67	48.67	40.89	10.03	10.70	11.07	10.60
Mean B	76.99	91.01	96.86		43.92	47.25	53.00		9.76	9.96	10.96	
CD at 5%	A=7.30	B=4.47	A+B=12.64		A=4.82	B=2.95	A+B=8.35		A=7.30	B=4.47	A+B=12.64	

Varieties (V): V1-VTLCH-1, V2- VTLCH-2, V3- VTLCH-3, V4- VTLCH-4, V5- VTLCH-5, V6- VTLCC-1, V7- VTLCS-1, V8-VTLCS-2
 N levels (N): N₁- N @ 40 ppm, N₂- N @ 80 ppm, N₃-N @ 120 ppm. SEM- Standard Error of means, CD- Critical difference

Table 2. Fresh weight of leaves, stem, root and whole plant of cocoa varieties as influenced by varied N levels

Varieties	Leaves				Stem				Root				Whole Plant			
	N1	N2	N3	Mean A	N1	N2	N3	Mean A	N1	N2	N3	Mean A	N1	N2	N3	Mean A
V1	64.45	67.68	94.64	75.59	53.65	56.74	79.24	63.21	64.45	29.48	18.63	37.52	182.55	153.89	192.52	176.32
V2	59.78	78.28	67.77	68.61	44.74	50.35	45.35	46.81	17.98	26.70	23.93	22.87	122.50	155.33	137.05	138.29
V3	45.89	66.95	72.49	61.78	33.97	48.42	54.79	45.73	25.03	30.36	24.20	26.53	104.90	145.72	151.47	134.03
V4	53.60	80.50	91.24	75.11	37.53	54.48	69.15	53.72	36.66	27.75	29.44	31.28	127.78	162.74	189.83	160.12
V5	57.51	61.92	86.15	68.53	35.41	36.78	69.22	47.14	15.65	23.79	30.54	23.33	108.57	122.49	185.92	138.99
V6	72.30	66.85	77.29	72.15	60.71	46.42	62.65	56.59	58.66	20.91	24.67	34.75	191.67	134.18	164.61	163.49
V7	68.46	84.60	62.61	71.89	57.08	58.49	48.52	54.70	31.20	30.49	22.21	27.97	156.74	173.58	133.34	154.55
V8	42.36	56.07	66.59	55.01	44.71	48.32	61.48	51.50	31.62	30.00	20.53	27.38	118.69	134.39	148.60	133.89
Mean B	58.04	70.36	77.35		45.98	50.00	61.30		35.16	27.43	24.27		139.17	147.79	162.92	
CD at 5%	A=5.34	B=3.27	A+B=9.26		A=4.75	B=2.91	A+B=8.23		A=3.22	B=1.97	A+B=5.58		A=11.22	B=6.87	A+B=19.43	

Varieties (V): V1-VTLCH-1, V2- VTLCH-2, V3- VTLCH-3, V4- VTLCH-4, V5- VTLCH-5, V6- VTLCC-1, V7- VTLCS-1, V8-VTLCS-2
 N levels (N): N₁- N @ 40 ppm, N₂- N @ 80 ppm, N₃-N @ 120 ppm. SEM- Standard Error of means, CD- Critical difference

Table 3. Shoot-to-root, Root-to-leaf and Shoot-to-fine root ratios of cocoa varieties as influenced by varied N levels

Varieties	Shoot:Root				Root:Leaf				Shoot:Fine root			
	N1	N2	N3	Mean A	N1	N2	N3	Mean A	N1	N2	N3	Mean A
V1	3.35	5.01	8.82	5.73	0.53	0.32	0.20	0.35	37.96	50.09	47.47	45.17
V2	6.12	4.94	5.11	5.39	0.28	0.32	0.30	0.30	61.19	50.64	46.16	52.66
V3	3.34	4.11	5.71	4.39	0.48	0.39	0.28	0.39	42.60	32.81	30.15	35.19
V4	2.83	5.15	5.93	4.63	0.60	0.31	0.28	0.40	61.08	55.14	44.36	53.53
V5	6.54	5.37	5.91	5.94	0.24	0.28	0.27	0.26	55.51	61.00	44.49	53.67
V6	3.12	5.81	5.83	4.92	0.53	0.27	0.28	0.36	31.24	36.48	42.35	36.69
V7	4.20	5.02	5.97	5.06	0.41	0.32	0.27	0.34	58.08	44.60	46.24	49.64
V8	2.93	3.78	5.71	4.14	0.58	0.44	0.32	0.45	42.78	30.63	42.94	38.78
Mean B	4.06	4.90	6.12		0.46	0.33	0.28		48.84	45.17	43.02	4.06
LSD at 5%	A=0.79	B=0.48	A+B=1.37		A=0.06	B=0.03	A+B=0.096		A=4.58	B=2.81	A+B=7.94	

Varieties (V): V1-VTLCH-1, V2- VTLCH-2, V3- VTLCH-3, V4- VTLCH-4, V5- VTLCH-5, V6- VTLCC-1, V7- VTLCS-1, V8-VTLCS-2
 N levels (N): N₁- N @ 40 ppm, N₂- N @ 80 ppm, N₃-N @ 120 ppm. SEM- Standard Error of means, CD- Critical difference

varieties. Increasing N levels significantly increased all the growth parameters. Cocoa hybrid, VTLCH-1 recorded the highest plant height (108.82 cm) and number of leaves (55.55), significantly higher than all other varieties. Among the interactions, plants of VTLCH-1 grown with a high N level of 120 ppm recorded the highest plant height (125.07 cm), number of leaves (69.33), and the highest stem diameter was observed in plants of VTLCH-5 grown with 120 ppm N (12.37 mm).

3.2 Biomass Production and Partitioning

Significant variations in biomass weight and partitioning were observed across different cocoa varieties, N levels, and their interactions. Higher N concentration in the nutrient solution significantly increased the fresh weight of leaves, stem, and the whole plant. However, root weight exhibited a declining trend with increasing N levels. Among the varieties, VTLCH-1 recorded the highest fresh weight of leaves (75.59 g), stem (63.21 g), roots (37.52 g), and the entire plant (176.32 g), significantly surpassing all other varieties. Among the interactions, VTLCH-1 grown at 120 ppm N recorded the highest values for fresh weight of leaves (94.64 g), stem (79.24 g), and the whole plant (192.52 g). In contrast,

the highest root weight (64.45 g) was observed in plants of VTLCH-1 grown with 40 ppm N (Table 2).

Significant variation in biomass partitioning to different parts was observed. Among different parts, biomass partitioning to leaves was high. At low nitrogen level, biomass partitioning to roots was high, whereas, at high nitrogen level, biomass partitioning to leaves was high. Among the varieties, VTLCH-5 exhibited the highest partitioning to leaves, VTLCH-1 showed the highest partitioning to stem and VTLCS-2 had the highest partitioning to roots (Fig. 1). Further, the biomass partitioning to different portions of the shoot was found to be significant. Among different parts of the shoot, the biomass partitioning to the middle shoot was high compared to the top and lower shoots. Among different N levels, biomass partitioning to top and lower shoots was highest at high N level, whereas the highest biomass partitioning to middle shoot was observed at medium N level. Among different varieties, VTLCH-2 recorded the highest biomass partitioning to the top shoot, VTLCH-3 recorded the highest biomass partitioning to the middle shoot and VTLCS-2 recorded the highest biomass partitioning to the lower shoot (Fig. 2).




Fig 1. Biomass partitioning to the leaf, stem and root of different cocoa varieties as influenced by varied N levels




Fig 2. Biomass partitioning to the top, middle and lower shoots of different cocoa varieties as influenced by varied N levels

Additionally, significant variation in biomass partitioning ratios was observed among the different cocoa varieties and nitrogen levels (Table 3). Cocoa hybrid, VTLCH-5 had the highest shoot-to-root ratio (5.94) and shoot-to-fine root ratio (53.67), while VTLCS-2 recorded the highest root-to-leaf ratio (0.45). Regarding nitrogen levels, plants grown with high N had the maximum shoot-to-root ratio (6.12), while those grown with low N had the highest root-to-leaf ratio (0.46) and shoot-to-fine root ratio (48.84). Considering the interaction between varieties and nitrogen levels, plants of VTLCH-1 grown with high nitrogen had the highest shoot-to-root ratio (8.82), while VTLCH-4 grown with low nitrogen had the lowest (2.83). In contrast, the highest root-to-leaf ratio was observed in VTLCH-4 grown with low nitrogen (0.60) and the lowest in VTLCH-1 grown with high nitrogen (0.20). The highest shoot-to-fine root ratio was recorded in VTLCH-2 grown with low nitrogen (61.19) and the lowest in VTLCH-3 grown with high nitrogen (30.15).

4. DISCUSSION

Nitrogen application encourages cell division, chlorophyll content, and photosynthesis, leading to increased biomass accumulation and more extensive foliage [9]. Additionally, the regulatory role of N in fundamental growth processes, including hormonal balance and enzyme activity,

significantly enhances growth parameters [10]. N helps in protein synthesis, a crucial factor in cell division and growth, resulting in increased biomass accumulation [11]. Ribeiro et al. [7] also reported increased biomass production with increasing N levels in cocoa.

Our findings align with those of Tian et al. [12] and Gaudin et al. [13], who observed enhanced root elongation at low nitrogen levels, suggesting that moderate nitrogen application supports root development. Conversely, Wang et al. [14] found that high nitrogen availability reduces root biomass, indicating that excess nitrogen may shift the allocation of resources away from root growth. Comfort et al. [15] further suggested that excessive nitrogen application can impede overall plant growth due to high nitrate levels in the buds, which inhibit starch synthesis and reduce root sugar levels, ultimately affecting root growth [16]. Zhang et al. [17] also noted that excess nitrogen promotes aboveground growth while decreasing root growth, resulting in a higher shoot-to-root ratio. Niu et al. [18] observed similar effects in citrus, where excessive nitrogen application increased aboveground biomass and decreased root growth.

These collective findings indicate that high nitrogen application positively impacts leaf and stem biomass production but hinders root biomass production in cocoa. This suggests that

high nitrogen application favors the allocation of biomass to leaves and stems at the expense of roots. Consequently, high nitrogen levels in cocoa are likely to increase the percentage of biomass allocated to leaf and stem growth while reducing the biomass allocated to roots. This shift in biomass partitioning highlights the importance of balancing nitrogen application to optimize overall plant development and maintain a more favorable shoot-to-root ratio.

5. CONCLUSION

This study highlights the significant effects of nitrogen (N) levels and varietal differences on growth, biomass production, and biomass partitioning in cocoa varieties. Different cocoa varieties responded distinctly to varying nitrogen levels, affecting overall growth and biomass allocation. VTLCH-1 recorded the highest plant height, number of leaves, and biomass production, particularly at the high N level. Higher nitrogen levels enhanced biomass production in leaves and stems across most varieties, while root biomass was adversely affected, leading to an increased shoot-to-root ratio. This shift in biomass partitioning underscores the necessity of optimizing nitrogen application and selecting suitable cocoa varieties for balanced growth and improved productivity.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

The authors thank the Director of ICAR-Central Plantation Crops Research Institute, Kasaragod, India. This research received funding from the Indian Council of Agricultural Research (ICAR-CPCRI Institute Project Code no. 1000763058) and KSTePS, Department of Science and Technology (DST), GOVT. OF KARNATAKA. KSTePS had no role in the study design, data collection, data analysis, decision to publish, or manuscript preparation

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Karpagalakshmi, Muthusamy A. A study on export performance of cocoa products in India. *International Journal of Innovative Research in Multidisciplinary Field*. 2019; 5(2):101-106.
2. Hartemink AE, Donald LS. Nutrient stocks, nutrient cycling, and soil changes in cocoa ecosystems: A review. *Advances in Agronomy*. Academic Press. 2005;227-253.
3. De Souza SR, De Souza ESOE, Louro-Berbara RL, Fernandes MS, Stark EM. Enzymes of Nitrogen Metabolism and Proteases Activity in Hairy Roots of Clover and Carrots, with and without Arbuscular Mycorrhizal Fungi. *Journal of Plant Nutrition*. 2007;30:185–1204.
4. Fageria NK, Baligar VC. Enhancing nitrogen use efficiency in crop plants. *Advances in Agronomy*. 2005;88:97-185.
5. Wessel M, Fertilizer requirements of cacao (*Theobroma cacao* L.) in South-Western Nigeria. in: *Communication 61*. Koninklijk Instituut voor de Tropen, Amsterdam. 1971;72-73.
6. Kaba JSS, Zerbe D, Zanotelli AA, Abunyewa M, Tagliavni. Uptake of nitrogen by cocoa (*Theobroma cacao* L.) trees derived from soil decomposition of gliricidia (*Gliricidia sepium* Jacq.) shoots. *Acta Horticulture*. 2018;1217:263–269
7. Ribeiro MAQ, Silva JO, Aitken WM, Machado RCR, Baligar VC. Nitrogen Use Efficiency in Cacao Genotypes. *Journal of Plant Nutrition*. 2008;31:239-249.
8. Nayana H, Bhat R, Prasad MBN, Bhavishya, Apsara SE, Ramesh SV, Venkatesh J. Effect of zinc levels on growth, chlorophyll content and biomass allocation of cocoa varieties. *Journal of Experimental Agriculture International*. 2024;46(7):754-760.
9. Barker AJM. The uptake of zinc and calcium from solution culture by zinc tolerant and non-tolerant *Silene maritima* in relation to calcium supply. *New Phytology* 1978;81:321-330.
10. Crozier A, Kamiya Y, Bishop G, Yokota T. Biosynthesis of hormones and elicitor molecules, in: *Biochemistry and Molecular Biology of Plants*, Buchanan BB. (Eds), American Society of Plant Physiologists, Rockville, Maryland, 1999;850-929.
11. Chen C, Zhong Q, Lin Z, Yu W, Wang M, Chen Z, You X. Screening tea varieties for

- nitrogen efficiency. *Journal of Plant Nutrition*. 2017;40:1797-1804.
12. Tian QY, Chen FJ, Liu JX, Zhang FS, Mi GH. Inhibition of maize root growth by high nitrate supply is correlated with reduced IAA levels in roots. *Plant Physiology*. 2008;165:942-951.
 13. Gaudin ACM, McClymont SA, Holmes BM, Lyons E, Raizada MN. Novel temporal, fine-scale and growth variation phenotypes in roots of adult-stage maize (*Zea mays* L.) in response to low nitrogen stress. *Plant Cell Environ*. 2011;34:2122-2137.
 14. Wang Y, Mi GH, Chen FJ, Zhang JH, Zhang FS. Response of root morphology to nitrate supply and its contribution to nitrogen accumulation in Maize. *Journal of Plant Nutrition*. 2005;27:2189-2202.
 15. Comfort SD, Malzer GL, Busch R. Nitrogen fertilization of spring wheat genotypes: influence on root growth and soil water depletion. *Agronomy Journal*. 1988;80:114-120.
 16. Scheible W, Lauerer M, Schulze ED, Caboche M, Stitt M. Accumulation of nitrate in the shoot acts as a signal to regulate shoot-root allocation in tobacco. *Plant Journal*. 1997;11:671-691.
 17. Zhang HZ, Khan A, Tan DKY, Luo HH. Rational water and nitrogen management improves root growth, increases yield and maintains water use efficiency of cotton under mulch drip irrigation. *Frontiers in Plant Science*. 2017;8:912.
 18. Niu R, Zhuang Y, Lali MN, Zhao L, Xie J, Xiong H, Wang Y, He X, Shi X, Zhang Y. Root Reduction Caused Directly or Indirectly by High Application of Nitrogen Fertilizer Was the Main Cause of the Decline in Biomass and Nitrogen Accumulation in Citrus Seedlings. *Plants*. 2024;13:938.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/120306>