



Genetic Diversity Studies using D^2 Statistics in Pearl Millet [*Pennisetum glaucum* (L.) R. Br.] Genotypes under Aridity

Komal Shekhawat^{a*}, Nemichand Sharma^{b^o} and P. C. Gupta^{c[#]}

^a Department of Genetics and Plant Breeding, College of Agriculture, Swami Keshwanand Rajasthan Agricultural University, Bikaner (Rajasthan)-334006, India.

^b Swami Keshwanand Rajasthan Agricultural University, Bikaner (Rajasthan)-334006, India.

^c Department of Genetics and Plant Breeding, Swami Keshwanand Rajasthan Agricultural University, Bikaner, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i930802

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84004>

Received 15 December 2021

Accepted 23 February 2022

Published 26 April 2022

Original Research Article

ABSTRACT

The present study was carried out to estimate the genetic diversity for yield and its contributing traits in pearl millet (*Pennisetum glaucum* (L.) R. Br.) Genotypes. The experimental material was consisting of 32 pearl millet released genotypes. The experiment was laid out in Randomized Block Design with three replications during *Kharif* 2021 under rain fed conditions. The observations were recorded on ten randomly selected plants from each genotype in each replication for nine different quantitative traits. Based on D^2 values, 32 genotypes were grouped into 3 clusters. Among 3 clusters, cluster I had more number of genotypes (23) followed by cluster II (6), and cluster III (3). The maximum intra cluster distance was observed in cluster III (12.2). The highest inter cluster distance was observed between cluster III and cluster I (18.29) followed by cluster III and cluster II (16.72), and cluster II and cluster I (14.24). The cluster III (49.96) had significant and higher cluster mean for seed yield per plant.

Keywords: Pearl millet; genetic diversity; intra cluster distance; inter cluster distance.

^{*}Ph. D. scholar;

^oSenior Research Fellow;

[#]Professor;

*Corresponding author: E-mail: komalshekhawat2506@gmail.com;

1. INTRODUCTION

Pearl millet commonly known as bulrush millet (*Pennisetum glaucum* L.). R. Br. also classified as *P. typhoides*, *P. americanum*, or *P. spicatum*, is a cultivated small-grain tropical cereal grass. Vernacular names include: “bajra” (India), “gero” (Nigeria, Hausa language), “hegni” (Niger, Djerma language), “sanyo” (Mali), “dukhon” (Sudan, Arabic), and “mahangu” (Namibia). Pearl millet is quantitatively the most important millet, with world annual production ~14 million tons (Mt). It is cultivated mainly in the semi-arid tropics, almost exclusively by subsistence and small-scale commercial farmers.

Grain yield is an unpredictable character and relies upon number of component characters which are quantitatively inherited. Thus, before starting any breeding programmes, intensive information on the nature and greatness of genetic variability is extremely essential. The presence of genetic variability is of utmost importance for any breeding programme and due to this reason the plant breeders have emphasized the evaluation of germplasm for the improvement of crop yield as well as for utilization in further breeding programmes [1-3]. Evaluation of plant genetic resources is a pre-requisite for which the future breeding work is based. Crop improvement depends on the magnitude of genetic variability present in the base population.

Genetic diversity is one of the criteria for parent's determination in the hybridization programmes. The accessibility of transgressive segregant in any breeding project depends upon the diversity between the parents associated. The evaluation of genetic diversity through biometrical methodology by Mahalanobis D^2 -statistic has made conceivable to pick genetically diversified parents [2,4]. The divergence analysis has a definite job to play in an effective decision of dissimilar parents for hybridization to exploit greatest heterosis or transgressive segregants for improving the grain yield. Identification of high grain yield traits of pearl millet genotypes is important to further breeding programmes to develop a hybrids or cultivar. This is done through variability and genetic divergence analysis.

2. MATERIALS AND METHODS

The present experiment was carried out at Agricultural Research Station, Bikaner during

Kharif, 2021. The research farm is situated between 27°1' N latitude and 71°54' E longitude at an altitude of 228.50 meters above mean sea level. This region falls under agro-climatic zone 1C of Rajasthan. The experimental material for present investigation comprised of thirty two released genotypes of pearl millet. The experimental field was divided into 3 blocks of equal size and each plot consisted of six rows each of 4 meter length with row spacing of 60 cm and plant to plant spacing of 15 cm. All recommended agronomical cultural practices were carried out to raise a good crop. Observations were recorded based on ten randomly selected plants in each genotype in each replication for all important characters viz., plant height (cm), no. of productive tillers/plant, panicle length (cm), panicle diameter (cm), test weight (g), dry fodder yield/plant (g), and grain yield/plant (g) while for days to 50% flowering and days to maturity, observations were recorded on plot basis.

2.1 Data Analysis

The analysis of replicated data was worked out to test the signification test. It was done according to the procedure of RBD for each character as per methodology suggested by Fisher [5]. The total variance and degree of freedom were partition into three components viz. replication, treatment and error. The data were subjected to analysis of variance adopting standard statistical methods. Estimation of genetic divergence is used in many plant breeding programs for parental selection. Mahalanobis's D^2 statistics and multivariate (cluster) analysis is a unique statistical tool for analysing genetic diversity. D^2 statistic was developed by Mahalanobis in [6]. Assessment of genetic diversity by using this technique in plant breeding was suggested by Rao[7].

2.2 Custer Analysis

The genotypes were grouped based on traits used in the primary analysis of cluster. The individuals with same descriptions are automatically grouped into same clusters. Euclidian or straight line measure is most commonly used statistic in diversity analysis based on biometrical traits was used. Euclidian distance among two individuals i and j having observations on traits denoted by x_{1k} , x_{2k} , x_{3k} x_{kk} computed by using formula given below:

$$d_{ij} = \sqrt{\sum_1^k (X_{ik} - X_{jk})^2}$$

Where, k = number of genotypes

2.3 Group Constellation

The genotypes are grouped into different clusters by using Tocher's method. The two genotypes having small inter-cluster distance from each other were considered as first, to which third genotype having small average D^2 value from the first two genotypes were added. Then the nearest fourth genotype added and it was done for remaining ones. At certain stage when it was felt that after adding particular genotype, the average D^2 value increase and this stage genotypes was not included in that cluster and another cluster was formed. Clusters were formed till all genotypes were added into one or other cluster.

2.4 Intra Cluster Distance

The intra-cluster D^2 was estimated by using the formulae $\sum D^2 i/n$, where $\sum D^2 i$ is the distance between all possible combinations [$n-1 (i-1)/2$] of the genotype (i) includes in a cluster.

2.5 Inter Cluster Distance

All possible D^2 values between the genotypes of two clusters were added with aid of $n_1 \times n_2$ for calculating inter-cluster distance. Where, n_1 and n_2 is number of genotypes in two clusters.

2.6 Cluster Means

Summing of mean values of all genotypes belonging to a particular cluster for each characters. It was divided by number of genotypes to obtain for cluster mean for a particular trait.

2.7 Character Contribution towards Divergence

All the possible combination of genotypes [$n (n-1)/2$], each trait was ranked based on d_i values ($y_i^1 - y_i^2$). The highest mean difference was rank first and lowest mean difference was rank "p", where "p" is total number of traits were studied.

Per cent contribution of traits = (Number of times appearing first in ranking by $X_i / 100 [\frac{n(n-1)}{2}]$)

Where, X_i = i^{th} character and n = number of genotypes.

3. RESULTS AND DISCUSSION

The mean sum of squares values for 9 biometrical traits was presented in Table-1. The mean sum of squares due to the genotypes were significant for all the characters studied at both level of significance 1% and 5%, suggesting the existence of high genetic variability among the genotypes for all the traits. This indicates that there is ample scope for selection of genotypes from the present gene pool for yield and its component traits. The presence of large amount of variability might be due to diverse source of material as well as environmental influence affecting the phenotypes.

3.1 Genetic Diversity

The mean value of grain yield and other components were converted into uncorrelated mean values. The D^2 values were calculated for all possible combinations through Tocher's [3] clustering method. Based on D^2 values, 32 genotypes were grouped into 3 clusters and are presented in Fig. 1 (Based on Dendrogram). Among 3 clusters, cluster I had more number of genotypes (23) followed by cluster II (6), and III (3).

3.2 Inter and Intra Cluster Distance

Inter and intra cluster distances are furnished in Table-2. The intra cluster values ranged from 7.25 to 12.2. The maximum intra cluster distance was observed in cluster III (12.2). The minimum intra cluster distance for cluster II (7.25). The inter cluster distance ranges from 14.24 (between I and II) to 18.29 (between I and III). Other inter cluster distance were between these values. The highest inter cluster distance was observed between cluster I and cluster III (18.29). Similar results were reported by Thakur et al. (2020) and Nimbalkar et al, [8].

3.3 Cluster Means

The results of cluster mean were furnished in Table-3, Cluster mean of days to 50% flowering ranged from 47.28 (cluster II) to 57.07 (cluster I). Similar results were reported by Thakur et al. (2020). Cluster mean of days to maturity were ranged from 78.11 (cluster II) to 87.75 (cluster I), similar results were reported by Jakhar et al. [9]. Cluster mean of plant height varied from 157.94 (cluster II) to 193.22 (cluster III). Cluster mean of productive tillers per plant varied from 1.54 (cluster I) to 2 (cluster III). Cluster mean of

Table 1. Mean sum of squares for different characters in pearl millet (*Pennisetum glaucum* (L.) R. Br.)

S. No.	Characters	Mean sum of squares		
		Replication (df=2)	Treatments (df=38)	Error (df=76)
1	Days to 50% flowering	8.573	76.366**	3.433
2	Days to 50% pod setting	10.51	82.688**	3.435
3	Plant height (cm)	67.885	1356.511**	22.423
4	Productive tillers/plant	0.03	0.446**	0.015
5	PL	2.698	11.923**	1.451
6	PD	0.066	0.163**	0.024
7	Test Weight	1.403	2.325**	0.527
8	Dry fodder yield/plant	71.651	1136.791**	56.209
9	Grain yield/plant	10.632	225.398**	4.64

**significant at 1% level

Table 2. Inter and intra cluster group distance of 32 genotypes of Pearl millet

Clusters	Cluster 1	Cluster 2	Cluster 3
Cluster 1	8.92	14.24	18.29
Cluster 2	14.24	7.25	16.72
Cluster 3	18.29	16.72	12.2




Fig. 1. Dendrogram for 9 biometrical traits towards genetic divergence

Table 3. Cluster means of 12 biometrical traits of 32 pearl millet genotypes

Genotypes	Days to 50% flowering	Days to maturity	Plant Height	Prod. tillers/plant	Panicle Length	Panicle Diameter	Test Weight	Dry fodder yield/plant	Grain yield/plant
Cluster 1	57.07	87.75	194.41	1.54	24.52	2.77	11.09	76.85	26.54
Cluster II	47.28	78.11	157.94	2.23	23.28	2.62	10.71	80.92	29
Cluster III	53.67	83.89	193.22	2	25.67	2.59	10.67	124.5	49.96




Fig. 2. % contribution of 12 biometrical traits towards genetic divergence

Table 4. % contribution of 9 biometrical traits of 32 pearl millet genotypes towards genetic divergence

	Traits	Times ranked first	Contribution %
1.	Days to 50% flowering	34	6.85%
2.	Days to maturity	1	0.20%
3.	Plant height	196	32.52%
4.	Productive tillers /plant	50	10.08%
5.	Panicle length	9	1.81%
6.	Panicle diameter	24	4.84%
7.	Test weight	6	1.21%
8.	Dry fodder yield/plant	30	6.05%
9.	Grain yield per plant	146	29.44%

Panicle Length were ranged from 23.28 (cluster II) to 25.67 (cluster III). Cluster mean of PD were reported from 2.59 (cluster III) to 2.77 (cluster I). Cluster mean of test weight ranged from 10.67 (cluster III) to 11.09 (cluster I). Cluster mean of dry fodder yield per plant were reported from 76.85 (cluster I) to 124.5 (cluster III). Cluster mean of grain yield per plant were reported from 26.54 (cluster I) to 49.96 (cluster III). Similar results were reported by Kuldeep et al. [10], Gaikwad et al. [11], Jakhar et al. [9], Nimbaldar et al. [8], Thakur et al. (2020), Jakhar et al. [9], Parashi et al. [12], Pandey et al. [13], Jivani et al. [14] and Thakur et al. (2020).

3.4 Contribution towards Divergence

The per cent contribution of 9 biometrical traits of 32 pearl millet genotypes towards genetic divergence was estimated and given in Table-4 and Fig. 2. The trait plant height (32.52%) had maximum contribution towards to genetic divergence followed by grain yield per plant (29.24%), productive tillers per plant (10.08%), days to 50% flowering (6.85%), dry fodder yield per plant (6.05%), Panicle Diameter(4.84%). Similar results were reported by Kudeep et al. [10], Jakhar et al. [9], Nimbalkar et al. [8], and Thakur et al. (2020).

4. CONCLUSION

It is concluded that maximum number of genotypes were grouped into 1st cluster which included 23 genotypes. The maximum intra cluster distance was shown by cluster III (12.2). The highest inter cluster distance was found between cluster III and cluster I (18.29). Therefore, genotypes present in these clusters may be used as parents to produce transgressive segregants.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ali M, Kumar S. An overview of pearl millet research in India. Indian J. Pulses Res. 2001;14(2):81-89.
2. Ibrkci H, Knewton SJB, Grusak MA. Pearl millet leaves as vegetable greens for humans: Evaluation of mineral composition. Journal of the Science of Food and Agriculture. 2003;83:945-950.
3. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. Indian Council of Agricultural Research. 1967; 103-108.
4. Varshney RK, Mohan SM, Gaur PM, Gangarao, NVPR, Pandey MK, Bohra A. Achievements and prospects of genomics-assisted breeding in three legume crops of the semi-arid tropics. Biotech Advance. 2013;31(8):1120–1134.
5. Fisher RA. The correlation between relative on the supposition of genotypes grown in Kumaun Himalaya. Indian Journal Genetics. 1936;66(1):37-38.
6. Mahalanobis PC. On the generalized distance in Statistics. Proceeding of the National Institute of Science of India. 1936;2: 49-55.
7. Rao CR. Advanced statistical methods in biometrical research. John Wiley and Sons Inc., New York. 1952;383.
8. Nimbalkar RD, Katre YY, Phad DS. Genetic diversity in Pearl millet (*Cicerarietinum* L.);2017.
9. Jakhar DS, Singh R, Kamble MS. Genetic diversity studies in pearl millet (*Cicerarietinum* L.) in Kolhapur region of

- Maharashtra. Bangladesh Journal of Botany. 2016;45(3):459-464.
10. Kuldeep R, Pandey S, Babbar A, Prakash V. Genetic diversity analysis in Pearl millet grown under heat stress conditions of Madhya Pradesh. Electronic Journal of Plant Breeding. 2015;6(4):962-971.
 11. Gaikwad AR, Desai NC, Pawar GH, Langhi AM. Genetic divergence in pearl millet. International Journal of Agricultural Sciences. 2014;10(2):812-815.
 12. Parashi VS, Lad TB, Mhase LB, Kute NS, Sonawane CJ. Genetic diversities studies in Pearl millet (*Cicerarietinum* L.). Bioinfolet. 2013;10(1B): 337-341.
 13. Pandey A, Gupta S, Kumar A, Thongbam PD, Pattanayak A. Genetic divergence, path coefficient and cluster analysis of pearl millet cultivars, in the mid altitudes of Meghalaya. Indian Journal of agricultural Sciences. 2013;83(12):1300-1304.
 14. Jivani JV, Mehta DR, Vaddoria MA, Raval L. Correlation and path coefficient analysis in pearl millet (*Cicer arietinum* L.). Research in Environment and Life Sciences. 2013;4(2):1167-1170.

© 2022 Shekhawat et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84004>